


Dietary fibre for the primary prevention of cardiovascular disease (Review)

Hartley L, May MD, Loveman E, Colquitt JL, Rees K



**THE COCHRANE
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2016, Issue 1

<http://www.thecochranelibrary.com>

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	3
OBJECTIVES	4
METHODS	4
Figure 1.	6
RESULTS	8
Figure 2.	10
Figure 3.	11
Figure 4.	12
DISCUSSION	13
AUTHORS' CONCLUSIONS	15
ACKNOWLEDGEMENTS	15
REFERENCES	15
CHARACTERISTICS OF STUDIES	20
DATA AND ANALYSES	56
Analysis 1.1. Comparison 1 Fibre versus control, Outcome 1 Total Cholesterol mmol/L change.	58
Analysis 1.2. Comparison 1 Fibre versus control, Outcome 2 HDL Cholesterol mmol/L change.	59
Analysis 1.3. Comparison 1 Fibre versus control, Outcome 3 LDL Cholesterol mmol/L change.	60
Analysis 1.4. Comparison 1 Fibre versus control, Outcome 4 Triglycerides mmol/L change.	61
Analysis 1.5. Comparison 1 Fibre versus control, Outcome 5 Systolic blood pressure (mmHg) change.	62
Analysis 1.6. Comparison 1 Fibre versus control, Outcome 6 Diastolic blood pressure (mmHg) change.	63
Analysis 2.1. Comparison 2 Subgroup analyses, Outcome 1 Total cholesterol mmol/L change.	64
Analysis 2.2. Comparison 2 Subgroup analyses, Outcome 2 HDL Cholesterol mmol/L change.	65
Analysis 2.3. Comparison 2 Subgroup analyses, Outcome 3 LDL Cholesterol mmol/L change.	67
Analysis 2.4. Comparison 2 Subgroup analyses, Outcome 4 Triglycerides mmol/L change.	68
Analysis 2.5. Comparison 2 Subgroup analyses, Outcome 5 Systolic blood pressure (mmHg) change.	69
Analysis 2.6. Comparison 2 Subgroup analyses, Outcome 6 Diastolic blood pressure (mmHg) change.	71
Analysis 2.7. Comparison 2 Subgroup analyses, Outcome 7 Total cholesterol mmol/L change.	72
Analysis 2.8. Comparison 2 Subgroup analyses, Outcome 8 HDL Cholesterol mmol/L change.	73
Analysis 2.9. Comparison 2 Subgroup analyses, Outcome 9 LDL Cholesterol mmol/L change.	75
Analysis 2.10. Comparison 2 Subgroup analyses, Outcome 10 Triglycerides mmol/L change.	76
Analysis 2.11. Comparison 2 Subgroup analyses, Outcome 11 Systolic blood pressure (mmHg) change.	78
Analysis 2.12. Comparison 2 Subgroup analyses, Outcome 12 Diastolic blood pressure (mmHg) change.	79
APPENDICES	79
CONTRIBUTIONS OF AUTHORS	86
DECLARATIONS OF INTEREST	86
SOURCES OF SUPPORT	86
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	87

[Intervention Review]

Dietary fibre for the primary prevention of cardiovascular disease

Louise Hartley¹, Michael D May², Emma Loveman³, Jill L Colquitt³, Karen Rees¹

¹Division of Health Sciences, Warwick Medical School, University of Warwick, Coventry, UK. ²Warwick Medical School, University of Warwick, Warwick, UK. ³Effective Evidence LLP, Eastleigh, UK

Contact address: Karen Rees, Division of Health Sciences, Warwick Medical School, University of Warwick, Coventry, Warwickshire, CV4 7AL, UK. Karen.Rees@warwick.ac.uk. rees_karen@yahoo.co.uk.

Editorial group: Cochrane Heart Group.

Publication status and date: New, published in Issue 1, 2016.

Review content assessed as up-to-date: 13 January 2015.

Citation: Hartley L, May MD, Loveman E, Colquitt JL, Rees K. Dietary fibre for the primary prevention of cardiovascular disease. *Cochrane Database of Systematic Reviews* 2016, Issue 1. Art. No.: CD011472. DOI: 10.1002/14651858.CD011472.pub2.

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

The prevention of cardiovascular disease (CVD) is a key public health priority. A number of dietary factors have been associated with modifying CVD risk factors. One such factor is dietary fibre which may have a beneficial association with CVD risk factors. There is a need to review the current evidence from randomised controlled trials (RCTs) in this area.

Objectives

The primary objective of this systematic review was to determine the effectiveness of dietary fibre for the primary prevention of CVD.

Search methods

We searched the Cochrane Central Register of Controlled Trials (CENTRAL) in *The Cochrane Library*, Ovid MEDLINE (1946 to January 2015), Ovid EMBASE (1947 to January 2015) and Science Citation Index Expanded (1970 to January 2015) as well as two clinical trial registers in January 2015. We also checked reference lists of relevant articles. No language restrictions were applied.

Selection criteria

We selected RCTs that assessed the effects of dietary fibre compared with no intervention or a minimal intervention on CVD and related risk factors. Participants included adults who are at risk of CVD or those from the general population.

Data collection and analysis

Two authors independently selected studies, extracted data and assessed risk of bias; a third author checked any differences. A different author checked analyses.

Main results

We included 23 RCTs (1513 participants randomised) examining the effect of dietary fibre. The risk of bias was unclear for most studies and studies had small sample sizes. Few studies had an intervention duration of longer than 12 weeks. There was a wide variety of fibre sources used, with little similarity between groups in the choice of intervention.

None of the studies reported on mortality (total or cardiovascular) or cardiovascular events. Results on lipids suggest there is a significant beneficial effect of increased fibre on total cholesterol levels (17 trials (20 comparisons), 1067 participants randomised, mean difference

-0.23 mmol/L, 95% CI -0.40 to -0.06), and LDL cholesterol levels (mean difference -0.14 mmol/L, 95% CI -0.22 to -0.06) but not on triglyceride levels (mean difference 0.00 mmol/L, 95% CI -0.04 to 0.05), and there was a very small but statistically significant decrease rather than increase in HDL levels with increased fibre intake (mean difference -0.03 mmol/L, 95% CI -0.06 to -0.01). Fewer studies (10 trials, 661 participants randomised) reported blood pressure outcomes where there is a significant effect of increased fibre consumption on diastolic blood pressure (mean difference -1.77 mmHg, 95% CI -2.61 to -0.92) whilst there is a reduction in systolic blood pressure with fibre but this does not reach statistical significance (mean difference -1.92 mmHg, 95% CI -4.02 to 0.19). There did not appear to be any subgroup effects by the nature of the type of intervention (fibre supplements or provision of foods/advice to increase fibre consumption) or the type of fibre (soluble/insoluble) although the number of studies contributing to each subgroup were small. All analyses need to be viewed with caution given the risks of bias observed for total cholesterol and the statistical heterogeneity observed for systolic blood pressure. Adverse events, where reported, appeared to mostly reflect mild to moderate gastrointestinal side-effects and these were generally reported more in the fibre intervention groups than the control groups.

Authors' conclusions

Studies were short term and therefore did not report on our primary outcomes, CVD clinical events. The pooled analyses for CVD risk factors suggest reductions in total cholesterol and LDL cholesterol with increased fibre intake, and reductions in diastolic blood pressure. There were no obvious effects of subgroup analyses by type of intervention or fibre type but the number of studies included in each of these analyses were small. Risk of bias was unclear in the majority of studies and high for some quality domains so results need to be interpreted cautiously. There is a need for longer term, well-conducted RCTs to determine the effects of fibre type (soluble versus insoluble) and administration (supplements versus foods) on CVD events and risk factors for the primary prevention of CVD.

PLAIN LANGUAGE SUMMARY

Dietary fibre to prevent cardiovascular disease

Background

Cardiovascular diseases (CVD) are a group of conditions affecting the heart and blood vessels. CVD is a global burden and varies between regions, and this variation has been linked in part to dietary factors. Such factors are important because they can be modified to help with CVD prevention and management. This review assessed the effectiveness of increased fibre intake as a supplement or in food stuffs in reducing cardiovascular death, all-cause death, non-fatal endpoints (such as heart attacks, strokes and angina) and CVD risk factors in healthy adults and adults at high risk of CVD.

Study characteristics

We searched scientific databases for randomised controlled trials (clinical trials where people are allocated at random to one of two or more treatments) looking at the effects of dietary fibre intake in healthy adults or those at high risk of developing CVD. We did not include people who already had CVD (e.g. heart attacks and strokes). The evidence is current to January 2015.

Key results

Twenty three trials fulfilled our inclusion criteria. All of the trials were short term and so could not examine the effect of fibre intake on CVD events. All of the trials examined the effects of fibre intake on lipid levels (lipids are fat-like substances, including cholesterol found in the blood), blood pressure or both. Pooling the results showed a beneficial reduction in total cholesterol and LDL cholesterol (sometimes called 'bad' cholesterol), and diastolic blood pressure with increasing fibre intake. There were no clear patterns for the type of fibre used (soluble or insoluble fibre) or the way in which fibre was provided (via supplements or food stuffs) but there were few studies in each group so results are uncertain.

Risk of bias of the included studies

Overall the risk of bias was unclear with few studies judged to be at low risk of bias (so less chance of arriving at the wrong conclusions because of favouritism by the participants or researchers), and for some there was a high risk of bias for some of the quality criteria. The results of this review need to be interpreted cautiously bearing this in mind. There is a need for longer-term well-conducted RCTs to determine the effects of fibre intake on CVD events and to further explore effects by the type of fibre and the way in which increased fibre is provided.

BACKGROUND

Description of the condition

Cardiovascular diseases (CVD) are a group of conditions that affect the heart and blood vessels and include coronary heart disease, cerebrovascular disease, and peripheral arterial disease (WHO 2013). One of the main mechanisms thought to cause CVD is atherosclerosis, where the arteries become clogged by atheromas or plaques (NHS 2012). CVD occurs when the arteries are completely blocked or when blood flow is restricted by a narrowed artery, limiting the amount of blood and oxygen delivered to organs or tissue (British Heart Foundation 2014). Arteries may naturally become harder and narrower with age, although this process may be accelerated by such factors as a sedentary lifestyle, obesity, ethnicity, smoking, high cholesterol, and high blood pressure (NHS 2012). Another cause of CVD is unstable plaque rupturing. It is thought that unstable plaques activate an inflammatory response in the body that causes the structure of atherosclerotic plaque to weaken and rupture, leading to the formation of blood clots (Spagnoli 2007).

CVD is the number-one cause of death and disability (WHO 2013) globally. Around 30% of total global deaths can be attributed to CVD (WHO 2013), and it is estimated to cause 17 million deaths per year (Bovet 2012). The World Health Organization (WHO) reports that by 2030, CVDs will account for almost 23.3 million deaths per year (WHO 2013). This burden is set to increase as a consequence of ageing populations and increasing levels of sedentary lifestyles and obesity.

One key public health priority in the prevention of CVD is targeting modifiable risk factors. One such risk factor is diet, which plays a major role in the aetiology of many chronic conditions, including CVD. Indeed, there are a number of dietary factors that have been found to be associated with a decrease in CVD risk, such as a low sodium intake (Aburto 2013), a low-carbohydrate diet (Hu 2014), intake of whole grains (Ye 2012), and a high consumption of fruits and vegetables (Begg 2007; Oude 2010). Such factors are important, not only because they have been linked to CVD development, but also because they can be modified. This makes them one of the main targets for interventions aimed at primary prevention and management of CVD.

Description of the intervention

To date, there is no globally accepted single definition for dietary fibre because of disagreements about which plant-derived substances should be included and how fibre values are derived (Buttriss 2008). In general, dietary fibre refers to the variety of plant substances that are resistant to the action of digestive enzymes (Eastwood 1983). Dietary fibre can be categorised into two

main groups: soluble and insoluble. Soluble fibre dissolves in water and delays the emptying of the stomach by forming a gel that slows digestion (Dietitians of Canada 2012). Sources of soluble fibre include bran, flaxseeds, oat cereal, and pears. Insoluble fibre, on the other hand, does not dissolve in water and speeds up the passage of food and waste through the stomach (Dietitians of Canada 2012). Sources of insoluble fibre include brown rice, barley, cabbage, celery, and whole grains. As each type of fibre aids the body in different ways, it is important that a healthy diet incorporates both soluble and insoluble fibre (NHS 2013).

Fibre consumption among the global population is low. For example, in the United Kingdom between 2008 and 2011, average intake of fibre was 12.8 g per day for women and 14.8 g per day for men (British Nutrition Foundation 2012). In the United States, the average fibre intake was 15.9 g per day for 2007 to 2008 (King 2012). Figures are similar in Japan and Malaysia (Nakaji 2002; Ng 1997). Current dietary recommendations for fibre intake range from 18 g per day (NHS 2013) to 40 g per day (King 2012; WHO 1990).

Little is known about the adverse effects of ingesting fibre over time (Bliss 2011), however, many studies have reported minor adverse events when administering gum arabic or psyllium in various doses to different populations (Jenkins 2002; Vuksan 2008). Indeed, Bliss 2011 found that when receiving fibre supplements, individuals with fecal incontinence experienced flatus, belching, fullness, and bloating.

How the intervention might work

The exact mechanisms by which dietary fibre reduces CVD risk are not known. However, when exposed to water, soluble fibre forms a gel in the stomach and small intestine that helps slow gastric emptying, hurry small intestine movement, and control nutrient absorption. In doing so, it is thought that soluble fibre reduces the effect of postprandial blood glucose and lipid increases (James 2003; Lunn 2007; Threapleton 2013), both of which are CVD risk factors. Furthermore, both soluble and insoluble fibre are thought to increase gastric distension and have an effect on gut hormones that increases satiety, leading to a lower food intake and, in the long term, weight reduction and improved glucose metabolism (Lattimer 2010; Satija 2012).

Dietary fibre has also been shown to increase the rate of bile acid excretion, which reduces total and low-density lipoprotein (LDL) cholesterol. In addition, once fermented in the colon, dietary fibre produces short-chain fatty acids that inhibit the synthesis of cholesterol (Lattimer 2010; Satija 2012). Finally, dietary fibre may have an impact upon plaque stability by decreasing pro-inflammatory cytokines known to affect plaque stability (Lattimer 2010). A recent meta-analysis of prospective cohort studies has shown a reduced risk of total mortality with increased fibre intake (pooled adjusted relative risk of total mortality for the highest category of dietary fibre intake versus the lowest was 0.77 (95% CI 0.74 to 0.8)

(Kim 2014). This has been confirmed in a recent observational cohort analysis of the PREDIMED trial (Buil-Cosiales 2014). A number of observational studies have also shown dietary fibre to have a beneficial association with CVD risk factors (Ascherio 1996; Eshak 2010; Kokubo 2011). One study showed an inverse relationship between dietary fibre and CVD risk in 39,876 female health professionals (Liu 2002), and another study showed that a high consumption of fibre is associated with a lower risk of incident ischaemic CVD in both men and women after a mean follow-up of 13.5 years (Wallström 2012). Further evidence on the beneficial association between dietary fibre and blood pressure and lipid levels can be found from systematic reviews of observational studies. Threapleton 2013 looked at evidence on dietary fibre and CVD risk from prospective cohort studies and found that a low risk of both CVD and coronary heart disease was associated with high dietary fibre intake.

Experimental studies have also shown dietary fibre to have a beneficial effect on CVD risk factors (Berg 2003; Saltzman 2001). For instance, Reyna-Villasmil 2007 found that oat-derived beta-glucan, when added to the American Heart Association Step 2 diet, improved the lipid profile of male participants with mild to moderate hypercholesterolaemia (Reyna-Villasmil 2007). In another study, ingesting oat cereal for six weeks was found to significantly reduce systolic blood pressure and diastolic blood pressure in hypertensive and hyperinsulinaemic participants when compared to a low-fibre cereal (Keenan 2002). Evidence also comes from systematic reviews of experimental studies. Streppel 2005 conducted a systematic review looking at dietary fibre and blood pressure and identified 24 relevant randomised controlled trials. The results from the meta-analyses showed that fibre supplementation caused a non significant reduction in systolic blood pressure but a significant reduction in diastolic blood pressure. Brown 1999 also conducted a systematic review examining dietary soluble fibre, but focused on blood cholesterol concentrations. They identified 67 controlled trials including 2990 participants that fulfilled their inclusion criteria and found that diets high in soluble fibre significantly reduced LDL and total cholesterol levels.

Why it is important to do this review

Few systematic reviews have been conducted that solely examine dietary fibre for CVD prevention. Those that have been carried out did not look at CVD events (Brown 1999; Streppel 2005; Whitehead 2014), involved limited searching (Streppel 2005), and did not assess the methodological rigour of their included studies (Brown 1999; Streppel 2005). With this in mind, we undertook this review to update current evidence on dietary fibre for the primary prevention of CVD by examining evidence from RCTs of dietary fibre in the general population as well as people at high risk of CVD. We included interventions of dietary advice to increase fibre consumption and the provision of high-fibre foods and fibre supplements.

OBJECTIVES

The primary objective of this systematic review was to determine the effectiveness of dietary fibre for the primary prevention of CVD.

METHODS

Criteria for considering studies for this review

Types of studies

Eligible studies were RCTs. We included studies reported as full text, those published as abstract only, and unpublished data.

Types of participants

Adults (age 18 and over) who are at high risk of CVD and adults from the general population in primary prevention trials were eligible. We excluded participants with the following co-morbidities/characteristics:

- those who have experienced a previous myocardial infarction (MI) or stroke, or both;
- those who have undergone a revascularisation procedure (coronary artery bypass grafting (CABG) or percutaneous transluminal coronary angioplasty (PTCA));
- those with angina or angiographically-defined coronary heart disease (CHD);
- those with type 2 diabetes, although this is a major risk factor for CVD, as interventions for the treatment and management of type 2 diabetes are covered by reviews registered with the Cochrane Metabolic and Endocrine Disorders Group.

Types of interventions

We included trials comparing dietary fibre with no intervention or minimal intervention (for example leaflets with no person-to-person intervention or reinforcement). The intervention was in the form of advice to increase consumption or the provision of fibre supplements or high-fibre foods. Where we found a sufficient number of trials, we stratified results by the type of fibre (soluble or insoluble), dose of supplementation, duration of intervention, and type of intervention (advice, diet, or supplementation).

We excluded multi-factorial lifestyle intervention trials and trials focused on weight loss in order to avoid confounding. We also focused on follow-up periods of 12 weeks (or three months) or more, as longer follow-up periods are more relevant for public health interventions.

Types of outcome measures

Primary outcomes

- All-cause mortality
- Cardiovascular mortality
- Non-fatal endpoints such as MI, CABG, PTCA, angina, angiographically-defined CHD, stroke, carotid endarterectomy, peripheral arterial disease (PAD)

Secondary outcomes

- Changes in blood pressure (systolic and diastolic) and blood lipids (total cholesterol, high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, triglycerides)
- Occurrence of type 2 diabetes as a major CVD risk factor
- Health-related quality of life
- Adverse effects
- Costs

Search methods for identification of studies

Electronic searches

We identified trials through systematic searches of the following bibliographic databases:

- The Cochrane Central Register of Controlled Trials (CENTRAL) (*The Cochrane Library* Issue 12, 2014)
- The Database of Abstracts of Reviews of Effects (DARE) (*The Cochrane Library* Issue 4, 2014)
- The NHS Economic Evaluation Database (NEED) (*The Cochrane Library* Issue 4, 2014)
- The Health Technology Assessment (HTA) Database (*The Cochrane Library* Issue 4, 2014)
- Ovid MEDLINE (1946 to January Week 1 2015)
- Ovid EMBASE and EMBASE Classic (1947 to 12 January 2015)

- Science Citation Index Expanded, Social Sciences Citation Index, and Conference Proceedings Citation Index - Science on Web of Science Core Collection (Thomson Reuters) (1970 to 12 January 2015)

We adapted the preliminary search strategy for MEDLINE (Ovid) ([Appendix 1](#)) for use in the other databases. We applied the Cochrane sensitivity-maximising RCT filter ([Lefebvre 2011](#)) to MEDLINE (Ovid) and adaptations of it to the other databases, except CENTRAL, DARE, NEED and HTA.

We also conducted a search of ClinicalTrials.gov (www.ClinicalTrials.gov) and the World Health Organization International Clinical Trials Registry Platform Search Portal (<http://apps.who.int/trialsearch/>).

We searched all databases from their inception to the present, and we imposed no restriction on language of publication.

Searching other resources


We checked reference lists of all primary studies and review articles for additional references. We also, where necessary, contacted authors for additional information.

Data collection and analysis

Selection of studies

Two authors (LH and MM) independently screened titles and abstracts for inclusion and coded them as 'retrieve' (eligible or potentially eligible/unclear) or 'do not retrieve'. We retrieved the full-text study reports/publications, and two authors (LH and MM) independently screened the full text to identify studies for inclusion, and identify and record reasons for exclusion of the ineligible studies. We resolved any disagreement through discussion or, where required, we consulted a third author (KR). We identified and excluded duplicates and collated multiple reports of the same study so that each study, rather than each report, was the unit of interest in the review. We recorded the selection process in sufficient detail to complete a PRISMA flow diagram ([Figure 1](#)) and [Characteristics of excluded studies](#) table ([Moher 2009](#)).

Figure 1. Study flow diagram.



Data extraction and management

We used a piloted data collection form for study characteristics and outcome data. Two authors (LH, MM or JC) extracted study characteristics from included studies. We extracted the following study characteristics.

- Methods: study design, total duration of study, details of any 'run in' period, number of study centres and location, study setting, withdrawals, and date of study.
- Participants: number (N), mean age, age range, gender, severity of condition, diagnostic criteria, inclusion criteria, and exclusion criteria.
- Interventions: intervention, comparison, concomitant medications, and excluded medications.
- Outcomes: primary and secondary outcomes specified and collected, and time points reported.

Two authors (LH, MM or JC) independently extracted outcome data from included studies. A third author (EL) resolved disagreements between the two reviewers. One author (EL) transferred data into a Review Manager ([RevMan 2014](#)) file. We double-checked that data was entered correctly by comparing the data presented in the systematic review with the study reports. A second author (KR) spot-checked study characteristics for accuracy against the trial report.

Assessment of risk of bias in included studies

Two authors (LH, MM or JC) independently assessed risk of bias for each study using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011a](#)). We resolved any disagreements by discussion or by involving another author (EL or KR). We assessed the risk of bias according to the following domains.

- Random sequence generation
- Allocation concealment
- Blinding of participants and personnel
- Blinding of outcome assessment
- Incomplete outcome data
- Selective outcome reporting
- Other bias (e.g. industry funding)

We graded each potential source of bias as high, low, or unclear and provide a quote from the study report together with a justification for our judgement in the Risk of bias section as part of the [Characteristics of included studies](#) table. We summarised the risk-of-bias judgements across different studies for each of the domains listed. Where information on risk of bias relates to unpublished data or correspondence with a trialist, we noted this in the Risk of bias section.

When considering treatment effects, we took into account the risk of bias for the studies that contribute to that outcome.

Assessment of bias in conducting the systematic review

We conducted the review according to the published protocol and report any deviations from it in the [Differences between protocol and review](#) section of the systematic review.

Measures of treatment effect

We analysed dichotomous data as odds ratios (OR) or risk ratios (RR) with 95% confidence intervals (CI) and continuous data as mean difference (MD) or standardised mean difference (SMD) with 95% CI. For continuous variables we presented data for the change from baseline rather than end-point data. We entered data presented as a scale with a consistent direction of effect, with the exception of HDL cholesterol where an increase in this outcome is a positive finding.

We narratively described skewed data reported as medians and interquartile ranges.

Unit of analysis issues

Studies with multiple intervention groups

In these cases, we used data from the control group for each intervention group comparison. We reduced the weight assigned to the control group by dividing the control group N by the number of intervention groups.

Cross-over trials

We included cross-over trials by using data only from the first half as a parallel group design. We only considered risk factor changes (for example, blood lipid levels and blood pressure) before participants crossed over to the other therapy and where the duration of intervention was a minimum of three months before cross-over.

Cluster randomised trials

We aimed to analyse cluster randomised trials by using the cluster (unit of randomisation) as the number of observations. Where needed, the individual level means and standard deviations adjusted for clustering would have been utilised together with the number of clusters in the denominator to appropriately weight the trials.

Dealing with missing data

We contacted investigators or study sponsors to verify key study characteristics and obtain missing numerical outcome data where possible (for example when a study is identified as abstract only). Where papers did not report results as change from baseline we calculated this and for the standard deviation differences followed the methods presented in the *Cochrane Handbook for Systematic Reviews of Interventions* for imputing these (16.1.3.2 Imputing standard deviations for changes from baseline Higgins 2011b), and assumed a correlation of 0.5 between baseline and follow-up measures as suggested by Follman 1992.

Assessment of heterogeneity

We used the I^2 statistic to measure heterogeneity among the trials in each analysis (Higgins 2003). Where we identified substantial heterogeneity (greater than 50%), we reported it and explored possible causes by pre-specified subgroup analysis.

Assessment of reporting biases

Where we were able to pool more than 10 trials, we created and examined a funnel plot to explore possible small-study biases for the primary outcomes (Sterne 2011).

Data synthesis

We undertook meta-analyses only where this was meaningful, that is if the treatments, participants, and the underlying clinical question were similar enough for pooling to make sense. Where there was no heterogeneity between included studies, we performed a fixed-effect meta-analysis. Where we detected substantial heterogeneity (I^2 greater than 50%) and could not explain it, we considered the following options: providing a narrative overview and not aggregating studies, or using a random-effects model with appropriately cautious interpretation.

Subgroup analysis and investigation of heterogeneity

We planned to carry out the following subgroup analyses.

- Type of fibre (soluble and insoluble)
- Dose of supplement
- Duration of intervention
- Type of intervention (fibre supplementation, provision of high-fibre foods, and advice to increase fibre consumption)

However, data were only available to undertake subgroup analyses on the type of fibre and the type of intervention.

We used the following outcomes in subgroup analyses.

- Changes in blood pressure (systolic and diastolic)
- Blood lipids (total cholesterol, HDL cholesterol, LDL cholesterol, triglycerides).

We used the formal test for subgroup interactions in Review Manager (RevMan 2014).

Sensitivity analysis

We planned to carry out sensitivity analyses looking at studies with a low risk of bias. However, no studies met this criteria (see [Risk of bias in included studies](#)). One study reported a high loss to follow up (> 20%) but this study did not report any data that could be included in the meta-analysis and therefore no sensitivity analysis was required.

Reaching conclusions

We based our conclusions only on findings from the quantitative or narrative synthesis of included studies for this review. We have avoided making recommendations for practice, and our 'Implications for research' suggests priorities for future research and outlines the remaining uncertainties in the area.

RESULTS

Description of studies

Results of the search

The searches generated 4207 hits after duplicates were removed. Screening of titles and abstracts identified 253 papers to go forward for formal inclusion and exclusion. Twenty three randomised controlled trials fulfilled the inclusion criteria and were included in the review. For a detailed description of the included studies see '[Characteristics of included studies](#)'. One ongoing study was also identified and is reported in [Characteristics of ongoing studies](#). One study is awaiting classification and is reported in [Characteristics of studies awaiting classification](#). The flow of studies through the review is presented in [Figure 1](#).

Included studies

Types of studies

Twenty three studies (in 27 publications) were included. Twenty were parallel RCTs and three were cross-over studies. One cross-over study was a cluster RCT (Nichenametla 2014), the remaining 22 trials were individually randomised. Of the studies, six were conducted in the USA. The other studies were conducted in Europe (Denmark, Finland, France, Italy, Norway, Spain, UK), Mexico, Japan, China or Australia. Only four studies reported the setting for the intervention, this was a university research clinic in

one (Perez-Jiminez 2008) and an outpatient setting another (Maki 2007) and two studies were set in the community (Nichenametla 2014; Pal 2011). Only one study reported the dates of the study (Tighe 2010 Wheat, being June 2005 to September 2008); the publication dates of all studies ranged from 1984 to 2014, with the majority being published after the year 2000.

The majority of trials were two-arm trials comparing the intervention with a placebo or control (18 trials). Of the remaining trials, four had three arms (in one study, Gato 2013, the third arm was irrelevant to this review) and one was a four-arm trial (although one arm was not relevant to this review, Pal 2011).

Overview of study populations

Trial sample sizes were generally small. Overall, 1513 participants were included in the trials. Six parallel trials and the one cluster RCT included study arms with 40 participants or more (Cicero 2010 Pysllium; He 2004; Lehtimaki 2005; Maki 2007; Nichenametla 2014; Pins 2002; Tighe 2010 Wheat). Other studies had fewer than 40 participants per study arm, with three studies including fewer than ten participants per arm (Aro 1984; Forcheron 2007; Hernandez-Gonzalez 2010).

Participants were described as 'healthy' in four studies (Forcheron 2007; He 2004; Jackson 1999; Marett 2004 Larch). In six studies participants were described as being overweight or obese (Birketvedt 2002; Hashizume 2012; Hernandez-Gonzalez 2010; Hu 2013; Pal 2011; Reimer 2013) and in three studies as having metabolic syndrome or signs of metabolic syndrome (Cicero 2010 Pysllium; Nichenametla 2014; Tighe 2010 Wheat). In three studies participants were described as having hypertension (Maki 2007; Pins 2002; Schlamovitz 1987). One study included women described as having climacteric symptoms (Makkonen 1993). The remaining six trials included participants with some degree of hypercholesterolaemia (Aro 1984; Gato 2013; Haskell 1992; Lehtimaki 2005; Perez-Jiminez 2008; Shimizu 2008).

Where reported, the mean age of participants was in the region of 35 years to 58 years in most studies. Two studies included participants between the ages of 19 years and 39 years (Forcheron 2007; Hu 2013) and one included participants aged 60 years to 61 years (Hashizume 2012). The proportion of male participants, where reported, was in the region of 32% to 73% with the exception of two studies, one which only included men (Aro 1984) and one which only included women (Makkonen 1993).

Description of interventions

In 15 studies the intervention was the provision of a fibre supplement. Eleven of these had a 12 week to 16 week follow-up, which was immediately at the end of the intervention in all except one study (Jackson 1999). Of these 15 studies, 14 had placebo comparisons, with the remaining study (Perez-Jiminez 2008) having a usual diet control group. Four studies had a six-month intervention with immediate follow-up. All except one study used a

placebo comparator; Cicero 2010 Pysllium had dietary advice as the comparator. The fibre supplements included in these studies varied. Supplements in 12 studies were judged to be soluble fibre (Aro 1984; Cicero 2010 Pysllium; Forcheron 2007; Haskell 1992; Hernandez-Gonzalez 2010; Jackson 1999; Lehtimaki 2005; Makkonen 1993; Marett 2004 Larch; Pal 2011; Reimer 2013; Schlamovitz 1987); in two studies insoluble fibre (Nichenametla 2014; Perez-Jiminez 2008) and in one study a combination of soluble and insoluble fibre (Birketvedt 2002).

In eight studies the intervention was the provision of foods high in fibre. All of these studies had an intervention and follow-up of 12 weeks duration. Of these studies five provided foods that were soluble fibre sources (Gato 2013; Hashizume 2012; He 2004; Maki 2007; Pins 2002); in two the fibre source was insoluble (Shimizu 2008; and one group of a three arm study Tighe 2010 Wheat); and in two a combination of soluble and insoluble fibre (Hu 2013; and one group of a three arm study, Tighe 2010 Wheat+Oats).

Outcomes included

None of the studies reported on mortality (total or cardiovascular) or cardiovascular events. In 22 studies lipid levels were outcomes and in twelve of these blood pressure was also reported as an outcome. One study (Maki 2007) only reported blood pressure as an outcome. Fourteen included studies reported adverse events, although there were limited details provided (Birketvedt 2002; Cicero 2010 GuarGum; Forcheron 2007; Hashizume 2012; Haskell 1992; He 2004; Hu 2013; Makkonen 1993; Marett 2004 Larch; Nichenametla 2014; Perez-Jiminez 2008; Pins 2002; Reimer 2013; Schlamovitz 1987).

No data for outcomes were reported in three studies (Gato 2013; Lehtimaki 2005; Maki 2007). One cross-over cluster RCT (Nichenametla 2014) did not report data for the first half of the trial and therefore we could not analyse data as specified in the review protocol.

One study is awaiting classification as the library were unable to locate it (Keenan 2002b). It is a costing study for one of the included studies (Pins 2002) and data on costs will be abstracted and included in the review when the paper becomes available.

Excluded studies

Details and reasons for exclusion for studies that closely missed the inclusion criteria are provided in the [Characteristics of excluded studies](#) table. Reasons for exclusion for the majority of studies included short term studies (< 3 months), the control group not being either no intervention or minimal intervention and alternative designs (not RCTs) (see [Figure 1](#)).


Risk of bias in included studies

A large proportion of studies were rated as unclear on many risk of bias domains (see Figure 2; Figure 3).

Figure 2. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Aro 1984	?	?	+	?	+	?	?
Birketvedt 2002	?	?	?	?	?	?	?
Cicero 2010 GuarGum	+	+	+	?	?	?	?
Cicero 2010 Pysllium	+	+	+	?	?	?	?
Forcheron 2007	?	?	+	?	+	?	?
Gato 2013	?	?	+	?	+	+	?
Hashizume 2012	?	?	+	?	?	?	?
Haskell 1992	?	?	+	?	?	?	?
He 2004	+	+	+	+	?	?	?
Hernandez-Gonzalez 2010	?	?	+	?	?	?	?
Hu 2013	?	?	?	?	+	?	?
Jackson 1999	?	?	+	?	+	?	?
Lehtimaki 2005	?	?	+	?	+	+	?
Maki 2007	?	?	+	?	+	+	?
Makkonen 1993	?	?	+	?	?	?	?
Marett 2004 Larch	?	?	+	?	?	+	?
Marett 2004 Tamarak	?	?	+	?	?	+	?
Nichenamella 2014	?	?	+	?	?	?	+
Pal 2011	?	?	+	?	+	?	?
Perez-Jiminez 2008	?	?	?	?	+	?	?
Pins 2002	?	?	+	+	+	?	+
Reimer 2013	?	?	+	?	?	?	?
Schlamovitz 1987	?	?	+	?	?	+	?
Shimizu 2008	?	?	+	?	+	?	?
Tighe 2010 Wheat	+	+	+	+	+	?	?
Tighe 2010 Wheat+Oats	+	+	+	+	+	?	?

Figure 3. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.



Allocation

Only three of 23 studies reported an adequate method of randomisation (Cicero 2010 Pysllium; He 2004; Tighe 2010 Wheat). These three studies also adequately reported the allocation concealment.

Blinding

Adequate blinding of participants and personnel were reported in 17 studies (Aro 1984; Forcheron 2007; Gato 2013; Hashizume 2012; Haskell 1992; He 2004; Hernandez-Gonzalez 2010; Jackson 1999; Lehtimaki 2005; Maki 2007; Makkonen 1993; Marett 2004 Larch; Nichenametla 2014; Pins 2002; Reimer 2013; Schlamovitz 1987; Shimizu 2008). In one study participants and personnel were unblinded (Cicero 2010 Pysllium) and in another the study was reported to be 'single' blind only (Pal 2011). In the remaining four studies blinding of participants and personnel was judged as unclear.

Adequate blinding of outcome assessors was reported in only two studies (He 2004; Tighe 2010 Wheat), the remaining studies were judged as unclear for detection bias.

Incomplete outcome data

Three studies did not have any missing data and were judged to be at low risk of attrition bias (Aro 1984; Gato 2013; Pins 2002).

Four studies reported different rates of drop outs or withdrawals between study groups (Lehtimaki 2005; Pal 2011; Perez-Jimenez 2008; Tighe 2010 Wheat); two studies reported drop outs or withdrawals for the total population but not per study group (Forcheron 2007; Shimizu 2008) and one study reported high loss to follow-up (Maki 2007). For the remaining studies this was not reported and judged at unclear risk of bias.

Selective reporting

Five studies did not report all outcomes as stated (Gato 2013; Lehtimaki 2005; Maki 2007; Marett 2004 Larch; Schlamovitz 1987), in three of these (Gato 2013; Lehtimaki 2005; Maki 2007) no data for any outcomes were reported.

Most other studies appeared to report all outcomes as intended, however, not enough information is available to check and these have been judged as unclear.

Other potential sources of bias

One cluster crossover randomised study (Nichenametla 2014) reported only two clusters and there was evidence of a carry-over effect, although we only intended to use data from the first half of the trial before crossover. One study was funded by the industry providing the cereals for the intervention and was judged to be

at high risk (Pins 2002). For the remaining studies this was not reported and judged at unclear risk of bias.

Effects of interventions

Three trials did not report data (Gato 2013; Lehtimaki 2005; Maki 2007), one trial did not report data that could be included in the pooled analysis (Pal 2011), and two cross-over studies (one also a cluster RCT) (Aro 1984; Nichenametla 2014) did not report data for the first half of the trial and therefore data could not be analysed as specified in the review protocol.

There were no data on the primary outcomes of the review.

Blood lipids


Total cholesterol. Eighteen studies reported total cholesterol and 17 of these could be summarised in a meta-analysis (20 com-

parisons) Analysis 1.1. The pooled analysis showed a significant difference between comparison groups where fibre decreased total cholesterol (MD -0.23 mmol/L, 95% CI -0.40 to -0.06, $P = 0.009$). There was significant heterogeneity ($P = 0.0001$; $I^2 = 65\%$) and results were pooled with a random-effects model.

One trial provided data that could not be included in the pooled analysis. Pal 2011 reported the percentage change in total cholesterol for their trial. Results showed that the fibre supplement (psyllium) group had a 21% reduction in total cholesterol compared to the control group ($P < 0.001$).

As no studies reported the review's primary outcomes a funnel plot was generated for total cholesterol as this analysis included the highest number of studies on which to assess publication bias. Inspection of the funnel plot suggests the possibility of publication bias (Figure 4).

Figure 4. Funnel plot of comparison: I Fibre versus control, outcome: I.I Total Cholesterol mmol/L change.



HDL cholesterol. Sixteen studies reported HDL cholesterol and 15 of these could be summarised in a meta-analysis (18 comparisons) Analysis 1.2. The pooled analysis showed a small but significant difference between comparison groups favouring the control

(an increase in HDL is beneficial and the direction of effect on the graphs have been changed for this outcome to illustrate this) (MD -0.03 mmol/L, 95% CI -0.06 to -0.01; $P = 0.02$). There was no

significant heterogeneity ($P = 0.70$; $I^2 = 0\%$). One trial provided data that could not be included in the pooled analysis. [Pal 2011](#) reported that there were no significant differences in the percentage change in HDL cholesterol between groups (fibre supplement, fibre supplement and dietary advice, and control) in their three-arm RCT.

LDL cholesterol. Sixteen studies reported LDL cholesterol and 15 of these could be summarised in a meta-analysis (18 comparisons) [Analysis 1.3](#). The pooled analysis showed a significant difference between comparison groups favouring fibre (MD -0.14 mmol/L, 95% CI -0.22 to -0.06; $P = 0.0006$). There was no significant heterogeneity ($P = 0.07$; $I^2 = 36\%$). One trial provided data that could not be included in the pooled analysis. [Pal 2011](#) reported the percentage change in LDL cholesterol. Results showed that the fibre supplement group had a 27% reduction in LDL cholesterol compared to the control group ($P < 0.001$).

Triglycerides. Sixteen studies reported triglycerides and 15 of these could be summarised in a meta-analysis (18 comparisons) [Analysis 1.4](#). The pooled analysis showed no significant difference between comparison groups (MD 0.00 mmol/L, 95% CI -0.04 to 0.05; $P = 0.88$). There was no significant heterogeneity ($P = 0.09$; $I^2 = 32\%$). One trial provided data that could not be included in the pooled analysis. [Pal 2011](#) reported the percentage change in triglycerides and showed that there was no significant difference between the fibre supplement and control group.

Blood pressure

Systolic blood pressure. Eight studies reported systolic blood pressure (SBP) and could be summarised in a meta-analysis (10 comparisons) [Analysis 1.5](#). The pooled analysis showed a reduction in SBP with the intervention but this did not reach statistical significance (MD -1.92 mmHg, 95% CI -4.02 to 0.19; $P = 0.07$). There was significant heterogeneity ($P = 0.0006$; $I^2 = 69\%$) and results were pooled with a random-effects model.

Diastolic blood pressure. Eight studies reported diastolic blood pressure and could be summarised in a meta-analysis (10 comparisons) [Analysis 1.6](#). The pooled analysis showed a significant difference between comparison groups favouring fibre (MD -1.77 mmHg, 95% CI -2.61 to -0.92; $P < 0.0001$). There was no significant heterogeneity ($P = 0.37$; $I^2 = 7\%$).

Subgroup analyses

Results are presented as subgroup analyses for the lipids and blood pressure outcomes for the type of intervention (fibre supplements or provision of foods high in fibre) and type of fibre (soluble, insoluble, combined fibre sources). Caution is required in the interpretation of some of these subgroup comparisons owing to low numbers of studies for some of these (described below).

Type of intervention

There did not appear to be any trends in the subgroup analyses of trials providing foods high in fibre or trials providing fibre supplements for lipids ([Analysis 2.1](#); [Analysis 2.2](#); [Analysis 2.3](#); [Analysis 2.4](#)). For blood pressure there were no significant subgroup effects seen for systolic blood pressure ([Analysis 2.5](#)) but there was a difference in effect between supplements and foods high in fibre for diastolic blood pressure in favour of providing foods high in fibre ([Analysis 2.6](#)) however this did not reach statistical significance ($P = 0.13$; $I^2 = 57.50\%$).

Type of fibre

In general it is difficult to establish any pattern in the study results for the subgroup looking at the source of fibre (soluble, insoluble or soluble and insoluble) owing to the small number of comparisons in the latter two subgroups. There were no statistically significant subgroup effects for lipids or systolic blood pressure. There was a difference in effect in fibre type for diastolic blood pressure with the largest effects seen for a combination of both soluble and insoluble fibre ([Analysis 2.12](#)), but this did not reach statistical significance ($P = 0.06$; $I^2 = 64.20\%$).

Adverse events

Fourteen trials reported information on adverse events. One study reported that no adverse effects of the interventions were observed ([Forcheron 2007](#)). One study reported that combined dietary (flatulence and diarrhoea) and antihypertensive medication side-effects scores decreased from baseline in the intervention but not the control group ([Pins 2002](#)). Four studies reported that participants had only a few gastrointestinal side-effects and that rates did not appear to differ between study groups ([Hu 2013](#); [Marett 2004](#) [Larch](#); [Reimer 2013](#); [Schlamovitz 1987](#)). Gastrointestinal side effects (predominantly flatulence but also constipation, nausea, bloating and diarrhoea) were more commonly reported in the fibre intervention groups than control groups in seven studies, although rates were generally low ([Birketvedt 2002](#); [Cicero 2010](#) [GuarGum](#); [Hashizume 2012](#); [Haskell 1992](#); [He 2004](#); [Makkonen 1993](#); [Perez-Jiminez 2008](#)). One cluster cross over trial only reported rates for the total population at the end of the intervention ([Nichenametla 2014](#)).

The remaining trials did not report on adverse events.

DISCUSSION

Summary of main results

This systematic review summarised 23 RCTs examining the effect of dietary fibre on risk factors for cardiovascular disease. None of the studies reported on mortality (total or cardiovascular) or cardiovascular events, the review's primary outcomes. Studies were

at risk of bias, few studies had an intervention duration of longer than 12 weeks and samples sizes were generally small. There was a wide variety of fibre sources used, with little similarity between groups in the choice of intervention.

Overall, there appears to be a significant effect of increased fibre on total cholesterol levels (MD -0.23 mmol/L, 95% CI -0.40 to -0.06), and LDL cholesterol levels (MD -0.14 mmol/L, 95% CI -0.22 to -0.06) but this effect is not demonstrated on triglyceride levels (MD 0.00 mmol/L, 95% CI -0.04 to 0.05), and there was a very small but statistically significant decrease rather than increase in HDL levels with increased fibre intake (MD -0.03 mmol/L, 95% CI -0.06 to -0.01). Fewer studies reported blood pressure outcomes where it appears there is a significant effect of increased fibre consumption on diastolic blood pressure (MD -1.77 mmHg, 95% CI -2.61 to -0.92) whilst there is a reduction in systolic blood pressure with fibre but this does not reach statistical significance (MD -1.92 mmHg, 95% CI -4.02 to 0.19). There did not appear to be any subgroup effects by the nature of the type of intervention (foods high in fibre versus fibre supplements) or the type of fibre (soluble or insoluble) although the number of studies contributing to each subgroup was small. All analyses need to be viewed with caution given the risks of bias of the studies, and for total cholesterol and systolic blood pressure, statistical heterogeneity was observed. Adverse events, where reported, appeared to mostly reflect mild to moderate gastrointestinal side-effects and these were generally reported more in the fibre intervention groups than the control groups.

Overall completeness and applicability of evidence

Whilst the number of trials meeting the inclusion criteria was relatively large, few studies had an intervention duration of longer than 12 weeks and samples sizes were generally small, so none reported on our primary outcomes, major CVD events.

There were a sufficient number of trials reporting our secondary outcomes, CVD risk factors, but heterogeneity between trials limited the findings, particularly for total cholesterol and systolic blood pressure where random-effects models were used to pool results. Heterogeneity between studies was due to the wide variety of fibre sources used and to differences in the participants recruited. We attempted to explore these differences in stratified analyses for fibre source (supplements versus provision of foods, and soluble versus insoluble fibre) where there were no obvious subgroup effects although the numbers of trials in each group were relatively small. There were insufficient trials to stratify results by cardiovascular risk.

Quality of the evidence

Trials were at risk of bias, with a large proportion of studies being rated as unclear on many quality criteria, and some studies at high risk of bias for individual quality domains.

The number of participants recruited was also generally small and so studies may be subject to small study bias (Nüesch 2010; Sterne 2000; Sterne 2001).

There was some evidence of publication bias from visual inspection of the funnel plot constructed for the outcome reported in most studies, total cholesterol [Figure 4](#).

Potential biases in the review process

The review authors carried out a comprehensive search across major databases for interventions involving dietary fibre for this review. In addition, the review authors screened the reference lists of systematic reviews and contacted study authors for information when needed. All screening, inclusion and exclusion and data abstraction were carried out independently by two review authors and analyses were conducted by one reviewer and checked by a second.

Multifactorial dietary interventions were excluded from this review because it would not be possible to disentangle the specific effects of fibre from other dietary interventions. We also excluded studies focusing on weight loss in order to reduce confounding. By restricting the comparison group to no intervention/placebo or minimal intervention we also reduced confounding. This did however limit the number of trials that were eligible for inclusion. We excluded a large number of trials of short duration (< 12 weeks) as we were interested in the sustained and longer-term effects of increased fibre intake, as these are more relevant for public health interventions.

Agreements and disagreements with other studies or reviews

The Global Burden of Disease Study conducted in 2010 found diets low in fibre to be one of the dietary risk factors for ischaemic heart disease (Lim 2012) and estimated that 11% of the disability-adjusted life years from ischaemic heart disease are attributable to diets low in fibre (Lim 2012). The primary source of this data is however from non-randomised studies. As demonstrated in the current systematic review, there is currently no RCT evidence on the effect of dietary fibre on mortality or heart disease.

There have been several previous systematic reviews examining the effects of increased fibre intake on cardiovascular risk factors on lipid levels (Brown 1999; Whitehead 2014) and blood pressure (Streppel 2005). These reviews used different inclusion criteria from the current review in terms of participants recruited (e.g. including patients with type 2 diabetes Whitehead 2014) and duration of interventions (e.g. including very short term studies Brown 1999), and were limited by searching (Streppel 2005) and lack

of assessment of methodological quality (Brown 1999; Streppel 2005) so the results are not directly comparable. Nevertheless, our review is in broad agreement with previous reviews in terms of the effects of dietary fibre on total cholesterol, LDL cholesterol (Brown 1999; Whitehead 2014) and blood pressure (Streppel 2005).

AUTHORS' CONCLUSIONS

Implications for practice

Currently there is no evidence from RCTs of the effects of dietary fibre on CVD clinical events. There is some evidence to suggest dietary fibre reduces total and LDL cholesterol and diastolic blood pressure but it is unclear which type of fibre (soluble or insoluble) and whether fibre supplements or provision of foods high in fibre are most effective and so no recommendations can be made for changing practice.

Implications for research

The majority of studies were short term and therefore did not report CVD clinical events, and it is unclear if the positive effects seen on total and LDL cholesterol and diastolic blood pressure are sustained over time. Studies were at risk of bias and there was insufficient information to determine the effects of fibre type and administration on outcomes. There is therefore a need for longer-term, well-conducted RCTs to determine the effects of fibre type (soluble versus insoluble) and administration (supplements versus foods) on CVD events and risk factors for the primary prevention of CVD.

ACKNOWLEDGEMENTS

We are grateful to Nicole Martin for conducting the searches for this review. With thanks also to Dr Frank Theis for providing additional data for their trial (Tighe 2010 Wheat; Tighe 2010 Wheat+Oats).

REFERENCES

References to studies included in this review

Aro 1984 *{published data only}*

Aro A, Uusitupa M, Voutilainen E, Korhonen T. Effects of guar gum in male subjects with hypercholesterolemia. *American Journal of Clinical Nutrition* 1984;**39**:911–6.

Birketvedt 2002 *{published data only}*

Birketvedt GS, Travis A, Langbakk B, Florholmen JR. Dietary supplementation with bean extract improves lipid profile in overweight and obese subjects. *Nutrition* 2002;**18**:729–33.

Cicero 2010 GuarGum *{published data only}*

Cicero AFG, Derosa G, Bove M, Imola F, Borghi C, Gaddi AV. Psyllium improves dyslipidaemia, hyperglycaemia and hypertension, while guar gum reduces body weight more rapidly in patients affected by metabolic syndrome following an AHA Step 2 diet. *Mediterranean Journal of Nutrition and Metabolism* 2010;**3**:47–54.

Cicero 2010 Psyllium *{published data only}*

Cicero AFG, Derosa G, Bove M, Imola F, Borghi C, Gaddi AV. Psyllium improves dyslipidaemia, hyperglycaemia and hypertension, while guar gum reduces body weight more rapidly in patients affected by metabolic syndrome following an AHA Step 2 diet. *Mediterranean Journal of Nutrition and Metabolism* 2010;**3**:47–54.

Forcheron 2007 *{published data only}*

Forcheron F, Beylot M. Long-term administration of inulin-type fructans has no significant lipid-lowering effect in normolipidemic humans. *Metabolism* 2007;**56**:1093–8.

Gato 2013 *{published data only}*

Gato N, Kadowaki A, Hashimoto N, Yokoyama S, Matsumoto K. Persimmon fruit tannin-rich fiber reduces cholesterol levels in humans. *Annals of Nutrition & Metabolism* 2013;**62**:1–6.

Hashizume 2012 *{published data only}*

Hashizume C, Kishimoto Y, Kanahori S, Yamamoto T, Okuma K, Yamamoto K. Improvement effect of resistant maltodextrin in humans with metabolic syndrome by continuous administration. *Journal of Nutritional Science and Vitaminology* 2012;**58**:423–30.

Haskell 1992 *{published data only}*

Haskell WL, Spiller GA, Jensen CD, Ellis BK, Gates JE. Role of water-soluble dietary fiber in the management of elevated plasma cholesterol in healthy subjects. *American Journal of Cardiology* 1992;**69**:433–9.

He 2004 *{published data only}*

Chen J, He J, Wildman RP, Reynolds K, Streiffer RH, Whelton PK. A randomized controlled trial of dietary fiber intake on serum lipids. *European Journal of Clinical Nutrition* 2006;**60**:62–8.

* He J, Streiffer RH, Muntner P, Krousel-Wood MA, Whelton PK. Effect of dietary fiber intake on blood pressure: a randomized, double-blind, placebo-controlled trial. *Journal of Hypertension* 2004;**22**:73–80.

Hernandez-Gonzalez 2010 *{published data only}*

Hernández-González SO, González-Ortiz M, Martínez-Abundis E, Robles-Cervantes JA. Chitosan improves insulin sensitivity as determined by the euglycemic-

- hyperinsulinemic clamp technique in obese subjects. *Nutrition Research* 2010;**30**:392–5.
- Hu 2013** *{published data only}*
Hu XJ, Gao JL, Zhang QY, Fu YQ, Li KL, Zhu SK, et al. Soy fiber improves weight loss and lipid profile in overweight and obese adults: a randomized controlled trial. *Molecular Nutrition & Food Research* 2013;**57**:2147–54.
- Jackson 1999** *{published data only}*
Jackson KG, Taylor GR, Clohessy AM, Williams CM. The effect of the daily intake of inulin on fasting lipid, insulin and glucose concentrations in middle-aged men and women. *British Journal of Nutrition* 1999;**82**:23–30.
- Lehtimäki 2005** *{published data only}*
Lehtimäki T, Metso S, Ylitalo R, Rontu R, Nikkila M, Wuolijoki E, et al. Microcrystalline chitosan is ineffective to decrease plasma lipids in both apolipoprotein E epsilon 4 carriers and non-carriers: a long-term placebo-controlled trial in hypercholesterolaemic volunteers. *Basic & Clinical Pharmacology & Toxicology* 2005;**97**:98–100.
- Maki 2007** *{published data only}*
Maki KC, Galant R, Samuel P, Tesser J, Witchger MS, Ribaya-Mercado JD, et al. Effects of consuming foods containing oat beta-glucan on blood pressure, carbohydrate metabolism and biomarkers of oxidative stress in men and women with elevated blood pressure. *European Journal of Clinical Nutrition* 2007;**61**:786–95.
- Makkonen 1993** *{published data only}*
Makkonen M, Simpanen AL, Saarikoski S, Uusitupa M, Penttilä I, Silvasti M, et al. Endocrine and metabolic effects of guar gum in menopausal women. *Gynecological Endocrinology* 1993;**7**:135–41.
- Marett 2004 Larch** *{published data only}*
Marett R, Slavin JL. No long-term benefits of supplementation with arabinogalactan on serum lipids and glucose. *Journal of the American Dietetic Association* 2004;**104**:636–9.
- Marett 2004 Tamarak** *{published data only}*
Marett R, Slavin JL. No long-term benefits of supplementation with arabinogalactan on serum lipids and glucose. *Journal of the American Dietetic Association* 2004;**104**:636–9.
- Nichenametla 2014** *{published data only}*
Nichenametla SN, Weidauer LA, Wey HE, Beare TM, Specker BL, Dey M. Resistant starch type 4-enriched diet lowered blood cholesterol and improved body composition in a double blind controlled cross-over intervention. *Molecular Nutrition & Food Research* 2014;**58**:1365–9.
- Pal 2011** *{published data only}*
* Pal S, Khossousi A, Binns C, Dhaliwal S, Ellis V. The effect of a fibre supplement compared to a healthy diet on body composition, lipids, glucose, insulin and other metabolic syndrome risk factors in overweight and obese individuals. *British Journal of Nutrition* 2011;**105**:90–100.
Pal S, Khossousi A, Binns C, Dhaliwal S, Radavelli-Bagatini S. The effects of 12-week psyllium fibre supplementation or healthy diet on blood pressure and arterial stiffness in overweight and obese individuals. *British Journal of Nutrition* 2012;**107**:725–34.
- Perez-Jimenez 2008** *{published data only}*
Perez-Jimenez J, Serrano J, Tabernerero M, Arranz S, Diaz-Rubio ME, Garcia-Diz L, et al. Effects of grape antioxidant dietary fiber in cardiovascular disease risk factors. *Nutrition* 2008;**24**:646–53.
- Pins 2002** *{published data only}*
Keenan JM, Pins JJ, Geleva D, Frazel C, O'Connor PJ, Cherney LM. Whole-grain oat cereal consumption reduces antihypertensive medication need: a cost analysis. *Preventive Medicine in Managed Care* 2002;**3**(1):9–17.
* Pins JJ, Geleva DRD, Keenan JM, Frazel C, O'Connor PJ, Cherney LM. Do whole-grain oat cereals reduce the need for antihypertensive medications and improve blood pressure control?. *The Journal of Family Practice* 2002;**51**(4):353–9.
- Reimer 2013** *{published data only}*
Reimer RA, Yamaguchi H, Eller LK, Lyon MR, Gahler RJ, Kacinik V, et al. Changes in visceral adiposity and serum cholesterol with a novel viscous polysaccharide in Japanese adults with abdominal obesity. *Obesity* 2013;**21**:E379–87.
- Schlamovitz 1987** *{published data only}*
Schlamowitz P, Halberg T, Warnoe O, Wilstrup F, Rytting K. Treatment of mild to moderate hypertension with dietary fibre. *Lancet* 1987;**2**:622–3.
- Shimizu 2008** *{published data only}*
Shimizu C, Kihara M, Aoe S, Araki S, Ito K, Hayashi K, et al. Effect of high beta-glucan barley on serum cholesterol concentrations and visceral fat area in Japanese men - A randomized, double-blinded, placebo-controlled trial. *Plant Foods for Human Nutrition* 2008;**63**:21–5.
- Tighe 2010 Wheat** *{published data only}*
Tighe P, Duthie G, Brittenden J, Vaughan N, Mutch W, Simpson WG, et al. Effects of wheat and oat-based whole grain foods on serum lipoprotein size and distribution in overweight middle aged people: a randomised controlled trial. *PLoS ONE* 2013; Vol. 8.
* Tighe P, Duthie G, Vaughan N, Brittenden J, Simpson WG, Duthie S, et al. Effect of increased consumption of whole-grain foods on blood pressure and other cardiovascular risk markers in healthy middle-aged persons: a randomized controlled trial. *American Journal of Clinical Nutrition* 2010;**92**:733–40.
- Tighe 2010 Wheat+Oats** *{published data only}*
Tighe P, Duthie G, Brittenden J, Vaughan N, Mutch W, Simpson WG, et al. Effects of wheat and oat-based whole grain foods on serum lipoprotein size and distribution in overweight middle aged people: a randomised controlled trial. *PLoS ONE* 2013; Vol. 8.
* Tighe P, Duthie G, Vaughan N, Brittenden J, Simpson WG, Duthie S, et al. Effect of increased consumption of whole-grain foods on blood pressure and other cardiovascular risk markers in healthy middle-aged persons: a randomized controlled trial. *American Journal of Clinical Nutrition* 2010;**92**:733–40.

References to studies excluded from this review

Bays 2013 *{published data only}*

Bays HE, Evans JL, Maki KC, Evans M, Maquet V, Cooper R, et al. Chitin-glucan fiber effects on oxidized low-density lipoprotein: a randomized controlled trial. *European Journal of Clinical Nutrition* 2013;**67**:2–7.

Davy 2002 *{published data only}*

Davy BM, Davy KP, Ho RC, Beske SD, Davrath LR, Melby CL. High-fiber oat cereal compared with wheat cereal consumption favorably alters LDL-cholesterol subclass and particle numbers in middle-aged and older men. *American Journal of Clinical Nutrition* 2002;**76**:351–8.

Dodson 1989 *{published data only}*

Dodson PM, Stephenson J, Dodson LJ, Kurnik D, Kritzinger EE, Taylor KG, et al. Randomized blind controlled trial of a high-fiber, low fat and low sodium dietary regimen in mild essential-hypertension. *Journal of Human Hypertension* 1989;**3**:197–202.

King 2008 *{published data only}*

King DE, Mainous AG, Egan BM, Woolson RF, Geesey ME. Effect of psyllium fiber supplementation on C-reactive protein: the trial to reduce inflammatory markers (TRIM). *Annals of Family Medicine* 2008;**6**:100–6.

Liao 2007 *{published data only}*

Liao FH, Shieh MJ, Chang NC, Chien YW. Chitosan supplementation lowers serum lipids and maintains normal calcium, magnesium, and iron status in hyperlipidemic patients. *Nutrition Research* 2007;**27**:146–51.

Mee 1997 *{published data only}*

Mee KA, Gee DL. Apple fiber and gum arabic lowers total and low-density lipoprotein cholesterol levels in men with mild hypercholesterolemia. *Journal of the American Dietetic Association* 1997;**97**:422–4.

Ramprasath 2014 *{published data only}*

Ramprasath VR, Jenkins DJA, Lamarche B, Kendall CWC, Faulkner D, Cermakova L, et al. Consumption of a dietary portfolio of cholesterol lowering foods improves blood lipids without affecting concentrations of fat soluble compounds. *Nutrition Journal* 2014;**13**:12.

Reid 2002 *{published data only}*

Reid R, Fodor G, Lydon-Hassen K, D'Angelo MS, McCrea J, Bowlby M, et al. Dietary counselling for dyslipidemia in primary care: results of a randomized trial. *Canadian Journal of Dietetic Practice and Research* 2002;**63**:169–75.

Tapola 2008 *{published data only}*

Tapola NS, Lyyra ML, Kolehmainen RM, Sarkkinen ES, Schauss AG. Safety aspects and cholesterol-lowering efficacy of chitosan tablets. *Journal of the American College of Nutrition* 2008;**27**:22–30.

References to studies awaiting assessment

Keenan 2002b *{published data only}*

Keenan JM, Pins JJ, Geleva D, Frazel C, O'Connor PJ, Cherney LM. Whole-grain oat cereal consumption

reduces antihypertensive medication need: a cost analysis. *Preventive Medicine in Managed Care* 2002;**3**(1):9–17.

References to ongoing studies

Lambert 2014 *{published data only}*

Lambert JE, Parnell JA, Han J, Sturzenegger T, Paul HA, Vogel HJ, et al. Evaluation of yellow pea fibre supplementation on weight loss and the gut microbiota: a randomized controlled trial. *BMC Gastroenterology* 2014;**14**:69. [<http://www.biomedcentral.com/1471-230X/14/69>]

Additional references

Aburto 2013

Aburto NJ, Ziolkovska A, Hooper L, Elliott P, Cappuccio FP, Meerpohl JJ. Effect of lower sodium intake on health: systematic review and meta-analyses. *BMJ* 2013;**346**:f1326.

Ascherio 1996

Ascherio A, Hennekens C, Willett WC, Sacks F, Rosner B, Manson J, et al. Prospective study of nutritional factors, blood pressure, and hypertension among US women. *Hypertension* 1996;**27**(5):1065–72.

Begg 2007

Begg S, Vos T, Barker B, Stevenson C, Stanley L, Lopez A. The burden of disease and injury in Australia 2003. Canberra: Australian Institute of Health and Welfare; 2007 May Vol. Cat. No. PHE 82.

Berg 2003

Berg A, Konig D, Deibert P, Grathwohl D, Berg A, Baumstark MW, et al. Effect of an oat bran enriched diet on the atherogenic lipid profile in patients with an increased coronary heart disease risk. A controlled randomized lifestyle intervention study. *Annals of Nutrition and Metabolism* 2003;**47**(6):306–11.

Bliss 2011

Bliss DZ, Savik K, Jung HJ, Whitebird R, Lowry A. Symptoms associated with dietary fiber supplementation over time in individuals with fecal incontinence. *Nursing Research and Practice* 2011;**60**(3 Suppl):S58–67.

Bovet 2012

Bovet P, Paccaud F. Cardiovascular disease and the changing face of global public health: a focus on low and middle income countries. *Public Health Reviews* 2012;**33**(2):397–415.

British Heart Foundation 2014

British Heart Foundation. Cardiovascular Disease 2014. <http://www.bhf.org.uk/heart-health/conditions/cardiovascular-disease.aspx> (accessed January 2014).

British Nutrition Foundation 2012

British Nutrition Foundation. Dietary Fibre 2012. <http://www.nutrition.org.uk/nutritionscience/nutrients/dietary-fibre?start=4> (accessed January 2014).

Brown 1999

Brown L, Rosner B, Willett WW, Sacks FM. Cholesterol-lowering effects of dietary fiber: a meta-analysis. *American Journal of Clinical Nutrition* 1999;**69**(1):30–42.

Buil-Cosiales 2014

Buil-Cosiales P, Itziar Zazpe I, Toledo E, Corella D, Salas-Salvadó J, Diez-Espino J, et al. Fiber intake and all-cause mortality in the Prevención con Dieta Mediterránea (PREDIMED) study. *American Journal of Clinical Nutrition* 2014;**100**(6):1498–507.

Buttriss 2008

Buttriss JL, Stokes CS. Dietary fibre and health: an overview. *Nutrition Bulletin* 2008;**33**:186–200.

Dietitians of Canada 2012

Dietitians of Canada. Food sources of soluble fiber. <http://www.dietitians.ca/getattachment/3bb6330f-0ab2-48fc-9d24-1303ad70003d/Factsheet-Food-Sources-of-Soluble-Fibre.pdf.aspx> (accessed 25 November 2013).

Eastwood 1983

Eastwood MA, Passmore R. Dietary Fiber. *Lancet* 1983;**2**:202–6.

Eshak 2010

Eshak ES, Iso H, Date C, Kikuchi S, Watanabe Y, Wada Y, et al. Dietary fiber intake is associated with reduced risk of mortality from cardiovascular disease among Japanese men and women. *Journal of Nutrition* 2010;**140**(8):1445–53.

Follman 1992

Follmann D, Elliott P, Suh I, Cutler J. Variance imputation for overviews of clinical trials with continuous response. *Journal of Clinical Epidemiology* 1992;**45**:769–73.

Higgins 2003

Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557–60.

Higgins 2011a

Higgins JPT, Altman DG, Sterne JAC (editors). Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Higgins 2011b

Higgins JPT, Deeks JJ, Altman DG (editors). Chapter 16: Special topics in statistics. In: Higgins JPT, Green S (editors), *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Hu 2014

Hu T, Bazzano LA. The low-carbohydrate diet and cardiovascular risk factors: evidence from epidemiologic studies. *Nutrition, Metabolism and Cardiovascular Diseases* 2014;**24**(4):337–43.

James 2003

James SL, Muir JG, Curtis SL, Gibson PR. Dietary fibre: a roughage guide. *Internal Medicine Journal* 2003;**33**:291–6.

Jenkins 2002

Jenkins DJ, Kendall CW, Vuksan V, Vidgen E, Parker T, Faulkner D, et al. Soluble fiber intake at a dose approved

by the US Food and Drug Administration for a claim of health benefits: serum lipid risk factors for cardiovascular disease assessed in a randomized controlled crossover trial. *American Journal of Clinical Nutrition* 2002;**75**(5):834–9.

Keenan 2002

Keenan JM, Pins JJ, Frazel C, Moran A, Turnquist L. Oat ingestion reduces systolic and diastolic blood pressure in patients with mild or borderline hypertension: a pilot trial. *The Journal of Family Practice* 2002;**51**(4):369.

Kim 2014

Kim Y, Je Y. Dietary fiber intake and total mortality: a meta-analysis of prospective cohort studies. *American Journal of Epidemiology* 2014;**180**:565–73.

King 2012

King DE, Mainous AG, Lambourne CA. Trends in dietary fiber intake in the United States, 1999–2008. *Journal of the Academy of Nutrition and Dietetics* 2012;**112**(5):642–8.

Kokubo 2011

Kokubo Y, Iso H, Saito I, Yamagishi K, Ishihara J, Inoue M, et al. Dietary fiber intake and risk of cardiovascular disease in the Japanese population: the Japan Public Health Center-based study cohort. *European Journal of Clinical Nutrition* 2011;**65**(11):1233–41.

Lattimer 2010

Lattimer JM, Haub MD. Effects of dietary fiber and its components on metabolic health. *Nutrients* 2010;**2**(12):1266–89.

Lefebvre 2011

Lefebvre C, Manheimer E, Glanville J. Lefebvre C, Manheimer E, Glanville J. Chapter 6: Searching for studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Lim 2012

Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;**380**(9859):2224–60.

Liu 2002

Liu S, Buring JE, Sesso HD, Rimm EB, Willett WC, Manson JE. A prospective study of dietary fiber intake and risk of cardiovascular disease among women. *Journal of the American College of Cardiology* 2002;**39**(1):49–56.

Lunn 2007

Lunn J, Buttriss JL. Carbohydrates and dietary fibre. *Nutrition Bulletin* 2007;**32**(1):21–64.

Moher 2009

Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Medicine* 2009;**6**(7):e1000097. [DOI: doi:10.1371/journal.pmed.1000097]

Nakaji 2002

Nakaji S, Sugawara K, Saito D, Yoshioka Y, MacAuley D, Bradley T, et al. Trends in dietary fiber intake in Japan over the last century. *European Journal of Nutrition* 2002;**41**(5): 222–7.

Ng 1997

Ng TKW. Dietary fat and fibre intakes of Malaysian adults: issues and implications when 'western targets' are set as dietary goals. *Malaysian Journal of Nutrition* 1997;**3**: 137–47.

NHS 2012

National Health Service. Atherosclerosis 2012. <http://www.nhs.uk/conditions/Atherosclerosis/Pages/Introduction.aspx#commentCountLink> (accessed January 2014).

NHS 2013

National Health Service. Why is fibre important? 2013. <http://www.nhs.uk/chq/pages/1141.aspx?categoryId=51> (accessed January 2014).

Nüesch 2010

Nüesch E, Trelle S, Reichenbach S, Rutjes AW, Tschannen B, Altman DG, et al. Small study effects in meta-analyses of osteoarthritis trials: meta-epidemiological study. *BMJ* 2010;**341**:c3515.

Oude 2010

Oude Griep LM, Geleijnse JM, Kronhout D, Ocke MC, Verschuern WM. Raw and processed fruit and vegetable consumption and 10-year coronary heart disease incidence in a population-based cohort study in the Netherlands. *PLoS ONE* 2010;**5**:e13609.

RevMan 2014 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Reyna-Villasmil 2007

Reyna-Villasmil N, Bermudez-Pirela V, Mengual-Moreno E, Arias N, Cano-Ponce C, Leal-Gonzalez E, et al. Oat-derived beta-glucan significantly improves HDLC and diminishes LDLC and non-HDL cholesterol in overweight individuals with mild hypercholesterolemia. *American Journal of Therapeutics* 2007;**14**:203–12.

Saltzman 2001

Saltzman E, Das SK, Lichtenstein AH, Dallal GE, Corrales A, Schaefer EJ, et al. An oat-containing hypocaloric diet reduces systolic blood pressure and improves lipid profile beyond effects of weight loss in men and women. *The Journal of Nutrition* 2001;**131**:1465–70.

Satija 2012

Satija A, Hu FB. Cardiovascular benefits of dietary fiber. *Current Atherosclerosis Reports* 2012;**14**(6):505–14.

Spagnoli 2007

Spagnoli LG, Bonanno E, Sangiorgi G, Mauriello A. Role of inflammation in atherosclerosis. *Journal of Nuclear Medicine* 2007;**48**(11):1800–15.

Sterne 2000

Sterne JA, Gavaghan D, Egger M. Publication and related bias in meta-analysis: power of statistical tests and prevalence in the literature. *Journal of Clinical Epidemiology* 2000;**53**:1119–29.

Sterne 2001

Sterne JA, Egger M, Smith GD. Systematic reviews in health care: investigating and dealing with publication and other biases in meta-analysis. *BMJ* 2001;**323**:101–5.

Sterne 2011

Sterne JAC, Egger M, Moher D (editors). Chapter 10: Addressing reporting biases. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Intervention*. Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Streppel 2005

Streppel MT, Arends LR, Van 't Veer P, Grobbee DE, Geleijnse JM. Dietary fiber and blood pressure: a meta-analysis of randomized placebo-controlled trials. *Archives of Internal Medicine* 2005;**165**:150–6.

Threapleton 2013

Threapleton DE, Greenwood DC, Evans CEL, Cleghorn CL, Nykjaer C, Woodhead C, et al. Dietary fibre intake and risk of cardiovascular disease: systematic review and meta-analysis. *BMJ* 2013;**347**:f6879.

Vuksan 2008

Vuksan V, Jenkins AL, Jenkins DJ, Rogovik AL, Sievenpiper JL, Jovanovski E. Using cereal to increase dietary fiber intake to the recommended level and the effect of fiber on bowel function in healthy persons consuming North American diets. *American Journal of Clinical Nutrition* 2008;**88**(5): 1256–62.

Wallström 2012

Wallström P, Sonestedt E, Hlebowicz J, Ericson U, Drake I, Persson M, et al. Dietary fiber and saturated fat intake associations with cardiovascular disease differ by sex in the Malmö Diet and Cancer Cohort: a prospective study. *PLoS One* 2012;**7**(2):e31637.

Whitehead 2014

Whitehead A, Beck EJ, Tosh S, Wolever TMS. Cholesterol-lowering effects of oat beta-glucan: A meta-analysis of randomized controlled trials. *American Journal of Clinical Nutrition*. 2014;**100**:1413–21.

WHO 1990

World Health Organization. *Diet, nutrition, and the prevention of chronic diseases. Report of a WHO Study Group, WHO Technical Report Series 797*. Geneva: World Health Organization, 1990.

WHO 2013

World Health Organization. Cardiovascular diseases (CVDs). Fact Sheet Number 317. March 2013. <http://www.who.int/mediacentre/factsheets/fs317/en/index.html> (accessed January 2014).

Ye 2012

Ye EQ, Chacko SA, Chou EL, Kugizaki M, Liu S. Greater whole-grain intake is associated with lower risk of type 2 diabetes, cardiovascular disease, and weight gain. *Journal of Nutrition* 2012;**142**(7):1304–13.

* *Indicates the major publication for the study*

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Aro 1984

Methods	<p>Setting: Finland, setting not reported Design: individual randomisation, cross-over RCT Dates: not reported Intervention duration: 12 weeks</p>	
Participants	<p>N: 14 randomised (7 in each group) Inclusion criteria: hypercholesterolaemia without marked hypertriglyceridaemia (serum cholesterol > 7.8 mmol/L and serum trig < 30 mmol/L), aged 30 years to 64 years Exclusion criteria: Thyroid dysfunction, renal or liver disease, diabetes, over 120% of ideal body weight Age (years): 51 years (all participants) Sex (% men): 100% Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): not reported Total cholesterol (mmol/L): intervention: 8.59 (0.7); control: 8.57 (0.8) HDL cholesterol (mmol/L): intervention: 1.49 (0.2); control: 1.47 (0.2) LDL cholesterol (mmol/L): intervention: 5.98 (0.7); control: 5.81 (0.6) Triglycerides (mmol/L): intervention: 1.64 (0.4); control: 1.88 (0.4) Blood pressure (mmHg): not reported</p>	
Interventions	<p>Type: fibre supplements Description: 15 g guar gum (soluble fibre), granulated guar gum preparation, 5 g doses at beginning of three main meals immediately after mixing with 200 ml of fluid. Control consumed 15 g placebo preparation consisting of wheat flour</p>	
Outcomes	Lipid levels, adverse events	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind with placebo

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing data
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	Unclear risk	Not enough information to judge

Birketvedt 2002

Methods	Setting: Norway, setting not reported Design: individual randomisation, parallel group Dates: not reported Intervention duration: three months	
Participants	N: 62 (intervention 31; control 31). Baseline characteristics and results based on Intervention 25 and control 27 Inclusion criteria: overweight and obese, 22 years to 66 years Exclusion criteria: not stated Age (years): mean (SD); intervention: 47 (11); control: 44 (11) Sex (% men): intervention: 36%; control: 44% Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): intervention: 33.9 (3.4); control: 35.0 (5.4) Total cholesterol (mmol/L): intervention: 6.6 (1.3); control: 6.2 (1.3) HDL cholesterol (mmol/L): intervention: 1.3 (0.5); control: 1.2 (0.4) LDL cholesterol (mmol/L): intervention: 4.4 (1.0); control: 4.1 (1.4) Triglycerides (mmol/L): intervention: 1.8 (0.8); control: 2.4 (2.2) Blood pressure (mmHg): not reported	
Interventions	Type: fibre supplements Description: extract of northern white kidney bean mixed with extract of locust bean gum (soluble and insoluble fibre). Two capsules three times each day at main meals. Control group were given placebo capsules, same regimen	
Outcomes	Lipid levels, adverse events	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement

Birketvedt 2002 (Continued)

Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No details reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Numbers and reasons given but no ITT
Selective reporting (reporting bias)	Unclear risk	Not enough information to judge
Other bias	Unclear risk	Not enough information to judge

Cicero 2010 GuarGum

Methods	<p>Setting: Italy, setting not reported Design: individual randomisation, parallel group Dates: not reported Intervention duration: six months</p>
Participants	<p>N: 137 randomised (intervention 1, N = 46, intervention 2, N = 46, control, N = 45) Inclusion criteria: metabolic syndrome, not taking antihypercholesterolaemic or antidiabetic treatments, or stabilised antihypercholesterol or antidiabetic treatment for at least six months. Any three of: waist circumference > 102 cm men and > 88 cm women; TG 1.70 mmol/L; HDL-C < 1.03 mmol/L men, < 1.29 women; SBP 130/85 mmHg or stabilised antihypertensive treatment for at least six months, FPG 6.1 mmol/L Exclusion criteria: severe hyperlipoproteinemias, uncontrolled diabetes, active treatments with anticoagulant, anticancer, antiretroviral drugs, previous CVD events Age (years): mean (SD); intervention 1: 58.4 (2.5); intervention 2: 57.9 (1.8); control: 57.2 (2.1) Sex (% men): not reported Ethnicity: 100% Caucasian Cardiovascular risk status: mean (SD) BMI (kg/m²): intervention 1: 28.8 (1.5); intervention 2: 28.6 (1.3); control: 28.5 (1.6) Total cholesterol (mmol/L): intervention 1: 4.8 (0.6); intervention 2: 4.9 (0.6); control: 4.8 (0.7) HDL cholesterol (mmol/L): intervention 1: 1.10 (0.13); intervention 2: 1.2 (0.1); control: 1.1 (0.1) LDL cholesterol (mmol/L): intervention 1: 2.9 (0.3); intervention 2: 3.0 (0.4); control: 3.0 (0.3) Triglycerides (mmol/L): intervention 1: 1.8 (0.3); intervention 2: 1.7 (0.2); control: 1.</p>

Cicero 2010 GuarGum (Continued)

	7 (0.3) Blood pressure (mmHg): intervention 1: SBP 131.4 (4.2); DBP 84.6 (4.4); intervention 2: 131.6 (4.0); DBP 85.2 (3.6); control: SBP 131.0 (3.7); DBP 84.5 (3.9)	
Interventions	Type: fibre supplement Description: intervention 1: soluble psyllium husk powder (soluble fibre); intervention 2: partially hydrolysed guar gum (soluble fibre). Both 3.5 g twice daily 20 minutes before two main meals, received a box containing a 60-day supply. Control group advised to follow the AHA Step 2 diet. All participants received dietary advice	
Outcomes	BP, lipid levels, adverse events	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomisation code by study statistician opening a previously prepared letter
Allocation concealment (selection bias)	Low risk	Code only broken after database lock
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not possible as two fibres have different colour and flavour and the comparison group was dietary advice not placebo
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details provided
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	ITT analysis but no other details
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	Unclear risk	Not enough information to judge

Cicero 2010 Pysllium

Methods	Please see details as above. This is the same trial with two relevant intervention arms for fibre supplement. intervention 1: soluble psyllium husk powder (Cicero 2010 Pysllium); intervention 2: partially hydrolysed guar gum (as above - Cicero 2010 GuarGum). Each arm was entered into the analysis separately.
Participants	

Cicero 2010 Pysllium (Continued)

Interventions		
Outcomes		
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	see previous
Allocation concealment (selection bias)	Low risk	see previous
Blinding of participants and personnel (performance bias) All outcomes	High risk	see previous
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	see previous
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	see previous
Selective reporting (reporting bias)	Unclear risk	see previous
Other bias	Unclear risk	see previous

Forcheron 2007

Methods	<p>Setting: France, setting not stated Design: individual randomisation, parallel design Dates: not reported Intervention duration: six months</p>
Participants	<p>N: 20 randomised (10 to each group). Baseline characteristics and results based on intervention N = 9, control N = 8 Inclusion criteria: Healthy, no history of diabetes, obesity or dyslipidaemia or taking any medications Exclusion criteria: unusual dietary habits or intensive physical activity Age (years): mean (SD); intervention: 29 (4); control: 34 (2) Sex (% men): intervention: 37.5%; control: 33.3% Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): not reported Total cholesterol (mmol/L): intervention: 4.48 (0.16); control: 3.91 (0.33)</p>

Forcheron 2007 (Continued)

	<p>HDL cholesterol (mmol/L): intervention: 1.29 (0.09); control: 1.03 (0.09) LDL cholesterol (mmol/L): intervention: 2.88 (0.13); control: 2.55 (0.33) Triglycerides (mmol/L): intervention: 0.71 (0.07); control: 0.78 (0.16) Blood pressure (mmHg): not reported</p>	
Interventions	<p>Type: fibre supplements Description: Fructans 5 g (soluble fibre) before breakfast and evening meal as powder dissolved in water, requested to avoid food rich in inulin or oligofructose, and to continue with their usual physical activity. Control participants had 5 g twice daily powder dissolved in water as a placebo. Same advice</p>	
Outcomes	Lipid levels, adverse events	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind with placebo
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	High risk	Not reported which groups or reasons for drop-outs
Selective reporting (reporting bias)	Unclear risk	All outcomes stated are reported but unable to fully judge
Other bias	Unclear risk	No details reported

Gato 2013

Methods	<p>Setting: Japan, setting not reported Design: individual randomisation, parallel group Dates: not reported Intervention duration: 12 weeks</p>
---------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Participants	<p>N: 27 randomised (intervention N = 13, control N = 14) Inclusion criteria: aged 25 years to 51 years, plasma cholesterol level 180-259 mg/dl Exclusion criteria: not stated Age (years): mean (SD); intervention: 36.4 (1.8); control: 36.6 (1.8) Sex (% men): intervention: 44%; control 40% Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): not reported Total cholesterol (mmol/L): not reported HDL cholesterol (mmol/L): not reported LDL cholesterol (mmol/L): not reported Triglycerides (mmol/L): not reported Blood pressure (mmHg): not reported</p>	
Interventions	<p>Type: fibre supplements Description: cookie bars containing 5 g of tannin-rich fibre (soluble fibre), three times a day before meals. Control group had cookie bars with 0 g of tannin-rich fibre. A low dose intervention group with 3 g of tannin-rich fibre was also reported but is not relevant here</p>	
Outcomes	Lipid levels	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind and placebo
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants reported
Selective reporting (reporting bias)	High risk	No data reported
Other bias	Unclear risk	Not enough detail to judge

Hashizume 2012

Methods	Setting: Japan, setting not reported Design: individual randomisation, parallel group Dates: not reported Intervention duration: 12 weeks	
Participants	N: 30 randomised (15 to each group). Inclusion criteria: mildly obese, exhibit visceral fat obesity and satisfy two of three from: dyslipidaemia, impaired glucose tolerance but no diabetes, high BP Exclusion criteria: DM, medical treatment with fibrates, thiazolidines or anion-exchange resins Age (years): mean (SD); intervention: 60.1 (8.9); control: 61.2 (11.6) Sex (% men): intervention: 60%; control: 73.3% Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): intervention: 28.1 (2.3); control: 26.8 (2.9) Total cholesterol (mmol/L): intervention: 5.74 (0.88); control: 5.64 (0.87) HDL cholesterol (mmol/L): intervention: 1.27 (0.24); control: 1.25 (0.33) LDL cholesterol (mmol/L): intervention: 3.81 (0.98); control: 3.77 (0.88) Triglycerides (mmol/L): intervention: 2.43 (0.46); control: 2.24 (0.50) Blood pressure (mmHg): not reported	
Interventions	Type: provision of foods high in fibre Description: unsweetened blended tea drink containing 9 g of resistant maltodextrins (RMD) (soluble fibre) in 280 ml. Other ingredients stated. Three bottles daily, one per meal time. Control: placebo tea, consumption as intervention group, nutrient contents same except fibre as placebo tea. Both groups not to change lifestyles with regards to diet, exercise or medication	
Outcomes	BP, lipid levels, adverse events	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind and placebo
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No details reported
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	Unclear risk	Not enough information to judge

Haskell 1992

Methods	<p>Setting: USA, setting not reported</p> <p>Design: individual randomisation, parallel group</p> <p>Dates: not reported</p> <p>Intervention duration: 12 weeks</p>
Participants	<p>N: 62 randomised (intervention N = 32, control N = 30). Baseline characteristics and results based on intervention N = 29, control N = 29</p> <p>Inclusion criteria: aged 20 years to 75 years, plasma cholesterol > 5.17 mmol/L, plasma triglycerides < 3.95 mmol/L, not on lipid lowering medication, dietary fibre supplementation, no history of gastrointestinal disease, fasting blood glucose < 6.6 mmol/L, normal liver function tests, no history or evidence of any disease that may influence lipoprotein metabolism, < 130% of ideal body weight, antihypertensive medications, oral contraceptions, oestrogen replacement if on a stable dose for at least six months and no plans to change</p> <p>Exclusion criteria: as above</p> <p>Age (years): not reported</p> <p>Sex (% men): not reported</p> <p>Ethnicity: not reported</p> <p>Cardiovascular risk status: mean (SD)</p> <p>BMI (kg/m²): not reported</p> <p>Total cholesterol (mmol/L): intervention: 5.97 (0.82); control: 6.05 (0.47)</p> <p>HDL cholesterol (mmol/L): intervention: 1.47 (0.47); control: 1.54 (0.44)</p> <p>LDL cholesterol (mmol/L): intervention: 4.01 (0.67); control: 3.86 (0.57)</p> <p>Triglycerides (mmol/L): intervention: 1.36 (0.61); control: 1.44 (0.64)</p> <p>Blood pressure (mmHg): not reported</p>
Interventions	<p>Type: fibre supplements</p> <p>Description: water soluble dietary fibre - 17 g/day from a mixture of dietary sources (major component being 56% acacia gum, the remaining psyllium husk and guar gum) (soluble fibre). Control given a placebo powder with only the carbohydrate fraction present, with equal calories</p>
Outcomes	Lipid levels, adverse events
Notes	
<i>Risk of bias</i>	

Haskell 1992 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind with placebo
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Numbers given but no reasons
Selective reporting (reporting bias)	Unclear risk	Not enough information to judge
Other bias	Unclear risk	Not enough information to judge

He 2004

Methods	<p>Setting: USA, setting not reported</p> <p>Design: individual randomisation, parallel group</p> <p>Dates: not reported</p> <p>Intervention duration: 12 weeks</p>
Participants	<p>N: 110 randomised (intervention 54, control 56).</p> <p>Inclusion criteria: healthy, systolic blood pressure 125 mmHg to 159 mmHg, diastolic blood pressure < 95 mmHg or DBP 75 mmHg to 94 mmHg with SBP < 160 mmHg; age 30 years to 65 years</p> <p>Exclusion criteria: current hypertension or use of antihypertensive medications, or those affecting blood pressure, hypercholesterolaemia or medications, diabetes or medications, severe obesity, history of CVD, history of chronic renal failure, serum creatinine at least 2.0 mg/dl, peptic ulcer, cancer, consumption of more than 21 alcoholic drinks per week, vegetarian, current use of fibre supplements, prior assignment to intervention in BP trial, poor compliance during run-in phase</p> <p>Age (years): mean (SD); intervention: 49.7 (8.2); control: 46.1 (8.6)</p> <p>Sex (% men): intervention: 40.7%; control: 39.3%</p> <p>Ethnicity: intervention: 57.9% African American; control: 53.6% African American</p> <p>Cardiovascular risk status: mean (SD)</p> <p>BMI (kg/m²): intervention: 28.5 (5.0); control: 29.3 (4.1)</p> <p>Total cholesterol (mmol/L): intervention: 5.12 (0.95); control: 5.18 (0.86)</p> <p>HDL cholesterol (mmol/L): intervention: 1.34 (0.43); control: 1.23 (0.36)</p> <p>LDL cholesterol (mmol/L): intervention: 3.19 (0.86); control: 3.32 (0.82)</p> <p>Triglycerides (mmol/L): intervention: 1.27 (0.73); control: 1.37 (0.63)</p>

He 2004 (Continued)

	Blood pressure (mmHg): intervention: SBP 129.8 (10.2); DBP 81.1 (6.4); control: SBP 126.4 (10.0); DBP 79.6 (6.6)	
Interventions	<p>Type: provision of foods high in fibre</p> <p>Description: daily serving of approximately 60 g of Quaker Oat Bran Concentrate as a muffin and 84 g of Quaker Oatmeal Squares (soluble fibre). Reduction in other high carbohydrate food intake. Remain at usual level of activity, alcohol intake and dietary sodium intake. Control group had 93 g of refined wheat as a muffin and 42 g of Kellogg's Corn Flakes</p>	
Outcomes	BP, lipid levels, adverse events	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Stratified by age and gender, block sizes reported
Allocation concealment (selection bias)	Low risk	Opaque envelopes only opened after confirmed eligibility
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind and cereals blinded
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Technicians masked to assignment
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Reported numbers but not reasons
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	Unclear risk	Not enough information to judge

Hernandez-Gonzalez 2010

Methods	<p>Setting: Mexico, setting not reported</p> <p>Design: individual randomisation, parallel group</p> <p>Dates: not reported</p> <p>Intervention duration: three months</p>
---------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Participants	<p>N: 12 randomised (six in each group)</p> <p>Inclusion criteria: nonsmokers, stable body weight, BP less than 130/80 mmHg, not using medication known to affect metabolism, BMI 30 kg/m² to 40 kg/m², 30 years to 50 years old</p> <p>Exclusion criteria: not reported</p> <p>Age (years): mean (SD); intervention: 41.6 (6.3); control: 42.6 (5.6)</p> <p>Sex (% men): intervention 50%; control 50%</p> <p>Ethnicity: not reported</p> <p>Cardiovascular risk status: mean (SD)</p> <p>BMI (kg/m²): intervention: 34.3 (2.7); control: 32.7 (1.7)</p> <p>Total cholesterol (mmol/L): intervention: 4.2 (1.0); control: 4.7 (0.8)</p> <p>HDL cholesterol (mmol/L): intervention: 0.9 (0.1); control: 0.9 (0.2)</p> <p>LDL cholesterol (mmol/L): intervention: 2.7 (0.7); control: 2.8 (0.7)</p> <p>Triglycerides (mmol/L): intervention: 2.4 (0.9); control: 2.2 (0.3)</p> <p>Blood pressure (mmHg): not reported</p>	
Interventions	<p>Type: fibre supplements</p> <p>Description: chitosam (soluble fibre) (750 mg, three times daily) 30 minutes before meals. All participants received general recommendations about their medical nutritional therapy and were instructed to not modify their usual forms of exercise. Control group received a placebo in the same pharmacological presentation as the intervention</p>	
Outcomes	Lipid levels	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Not stated
Selective reporting (reporting bias)	Unclear risk	Not enough information to judge

Other bias	Unclear risk	Not enough information to judge
------------	--------------	---------------------------------

Hu 2013

Methods	Setting: China, setting not reported Design: individual randomisation, parallel groups Dates: not reported Intervention duration: 12 weeks	
Participants	N: 39 randomised (intervention, N = 22; control, N = 17) Inclusion criteria: overweight and obese. Generally healthy adults aged 19 years to 39 years with BMI 23 to 35 Exclusion criteria: people taking any drugs, botanicals, or nutraceutical known to influence body weight or blood pressure, taking steroids and other agents that may influence lipid metabolism were excluded. Significant history of eating disorders, smoking and drinking, strenuous exercise to lose weight, hyperlipidaemia, hyperglycaemia, diabetes mellitus, cardiovascular disease, major systemic diseases, gastrointestinal problems, liver and renal failure Age (years): mean (SD) intervention: 24.7 (4.7); control: 21.2 (1.7) Sex (% men): intervention: 47%; control: 41% Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): intervention: 25.7 (1.9); control: 26.2 (2.5) Total cholesterol (mmol/L): intervention: 4.94 (0.9); control: 4.57 (1.0) HDL cholesterol (mmol/L): intervention: 1.35 (0.3); control: 1.28 (0.3) LDL cholesterol (mmol/L): intervention: 3.38 (1.83); control: 2.74 (3.6) Triglycerides (mmol/L): intervention: 0.94, (0.6); control: 1.24 (0.9) Blood pressure (mmHg): intervention: SBP 114.6 (11.7); DBP 75.2 (8.4); control: SBP 117.5 (13.2); DBP 74.0 (9.1)	
Interventions	Type: foods high in fibre Description: soy fibre (soluble and insoluble fibre) provided as biscuits. Average prescribed total dietary fibre intake was 27.5 g/day. Nutrient components of biscuit provided. 250 ml skim milk provided. Control group had control biscuits. Average prescribed total dietary fibre intake was 3.2 g/day. Soy fibre was substituted with flour-based ingredients. Approximately equivalent nutrient intake to intervention. Nutritional components of biscuit provided. 250 ml skim milk provided	
Outcomes	BP, lipid levels, adverse events	
Notes	It appears that more than 39 were randomised, as describes two withdrawing, but exact number randomised not given: 39 were analysed	
Risk of bias		
Bias	Authors' judgement	Support for judgement

Hu 2013 (Continued)

Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No details reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants included in analysis
Selective reporting (reporting bias)	Unclear risk	Data not fully reported for adverse events
Other bias	Unclear risk	Not enough information to judge

Jackson 1999

Methods	<p>Setting: UK, setting not reported Design: individual randomisation, parallel group Dates: not reported Intervention duration: eight weeks with twelve week follow-up</p>
Participants	<p>N: 54 randomised (27 in each group); no drop-outs Inclusion criteria: healthy, men and women, moderately raised fasting TC and TAG levels, aged between 35 years and 65 years, BMI 20 kg/m² to 32kg/m², no history of heart disease, diabetes or other endocrine or liver disease. Aerobic exercise fewer than three sessions of 30 mins/week. Not taking medications for hyperlipidaemia or that could affect plasma lipid levels, not trying to lose weight, taking antibiotics or dietary fatty acid supplements Exclusion criteria: covered above Age (years): mean (SD) intervention: 52.6 (8.6); control: 51.9 (10.5) Sex (% men): not reported Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): intervention: 26.5 (3.6); control: 26.1 (2.8) Total cholesterol (mmol/L): intervention: 5.86 (1.00); control: 6.43 (0.79) HDL cholesterol (mmol/L): intervention: 1.24 (0.28); control: 1.26 (0.28) LDL cholesterol (mmol/L): intervention: 3.97 (0.86); control: 4.55 (0.92) Triglycerides (mmol/L): intervention: 1.46 (0.55); control: 1.40 (0.40) Blood pressure (mmHg): not reported</p>

Jackson 1999 (Continued)

Interventions	<p>Type: fibre supplements Description: 10 g Inulin (soluble fibre) sachets, two mixed with water or other drinks. Provided as four-week batches. Control group consumed maltodextrins (placebo), provided and used in same way as intervention</p>	
Outcomes	Lipid levels	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details of randomisation schedule
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Low risk	No loss to follow-up
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	Unclear risk	Not enough information to judge

Lehtimaki 2005

Methods	<p>Setting: Finland, setting not reported Design: Individual randomisation, cross-over RCT Dates: not reported Intervention duration: three months (until first cross-over), seven-month study in total</p>
Participants	<p>N: 96 randomised (48 to each group) Inclusion criteria: middle aged men and women, plasma total cholesterol of 4.8 - 6.8 mmol/L with plasma triglyceride less than 3.0 mmol/L at end of one month run in. Aged 18-65 years Exclusion criteria: diabetes, renal, liver, adrenal disease, thyroid dysfunction, coronary artery or cerebrovascular disease, malignancy or terminal condition; use of lipid lowering</p>

	<p>medications, functional foods, or steroids; alcoholism or narcotic addiction; mental lability; pregnant, lactation and childbearing potential for women not using birth control, history of participation in another drug evaluation study within one month, severe adverse events caused by chitosan, allergy to crustaceans</p> <p>Age (years): reported for subgroups by genotype (apolipoprotein E ε4 carriers or not) only</p> <p>Sex (% men): reported for subgroups by genotype (apolipoprotein E ε4 carriers or not) only</p> <p>Ethnicity: not reported</p> <p>Cardiovascular risk status:</p> <p>BMI (kg/m²): reported for subgroups by genotype (apolipoprotein E ε4 carriers or not) only</p> <p>Total cholesterol (mmol/L): reported for subgroups by genotype (apolipoprotein E ε4 carriers or not) only</p> <p>HDL cholesterol (mmol/L): reported for subgroups by genotype (apolipoprotein E ε4 carriers or not) only</p> <p>LDL cholesterol (mmol/L): reported for subgroups by genotype (apolipoprotein E ε4 carriers or not) only</p> <p>Triglycerides (mmol/L): reported for subgroups by genotype (apolipoprotein E ε4 carriers or not) only</p> <p>Blood pressure (mmHg): reported for subgroups by genotype (apolipoprotein E ε4 carriers or not) only</p>	
Interventions	<p>Type: fibre supplements</p> <p>Description: hard gelatine capsules containing 400 mg of chitosan powder (soluble fibre). Three capsules twice a day before meals at 06:00 to 09:00 and 17:00 to 20:00. Control consumed placebo capsules with 400 mg starch, taken with same procedure</p>	
Outcomes	Lipid levels	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind and placebo, pharmacist numbered treatment containers to a randomisation list and codes not open until close of study
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported

Lehtimaki 2005 (Continued)

Incomplete outcome data (attrition bias) All outcomes	High risk	Numbers and details provided, difference in drop-outs between groups
Selective reporting (reporting bias)	High risk	No data reported
Other bias	Unclear risk	Not enough information to judge

Maki 2007

Methods	<p>Setting: USA, outpatient setting</p> <p>Design: individual randomisation, parallel group</p> <p>Dates: not reported</p> <p>Intervention duration: 12 weeks</p>
Participants	<p>N: 97 randomised (intervention N = 49; control N = 48); baseline characteristics and results based on intervention N = 26; control N = 34</p> <p>Inclusion criteria: elevated systolic blood pressure (130 mmHg to 179 mmHg) or diastolic blood pressure (85 mmHg to 109 mmHg) that was untreated or suboptimally controlled with one or two antihypertensive medications, or both. Waist circumference 87 cm or more for women and 95 cm or more for men and willing to maintain current weight and activity level during study period. Age 40 years or more</p> <p>Exclusion criteria: evidence of secondary cause of hypertension, use of over-the-counter drugs containing pseudoephedrine or other vasoconstrictors, mid-upper arm circumference 42 cm or more, daily fibre intake 20 g/day or more, pregnant, lactating, planning pregnancy or of childbearing potential not using approved contraception, use of omega-3 fatty acid or herbal supplements, weight-loss drugs, systemic corticosteroids, androgens, phenytoin, erythromycin, drugs other than stable-dose aspirin for regulating haemostasis, severe asthma, conditions that require periodic use of steroids, recent history or current diagnosis of unstable angina, CHF, MI, revascularisation, hypothyroidism, significant anaemia hyperadrenocorticalism, significant gastrointestinal, renal, pulmonary, hepatic or biliary disease, diabetes, cancer within five years, recent history or strong potential for alcohol or substance abuse</p> <p>Age (years): mean (SD); intervention: 63.0 (9.2); control: 57.1 (9.3)</p> <p>Sex (% men): intervention: 53.8%; control: 55.9%</p> <p>Ethnicity: intervention: white 69.2%, black 26.9%, other 3.8%; control: White 55.9%, Black 44.1, Other 0.0</p> <p>Cardiovascular risk status: mean (SD)</p> <p>BMI (kg/m²): intervention: 32.6 (5.1); control: 32.2 (5.3)</p> <p>Total cholesterol (mmol/L): not reported</p> <p>HDL cholesterol (mmol/L): not reported</p> <p>LDL cholesterol (mmol/L): not reported</p> <p>Triglycerides (mmol/L): not reported</p> <p>Blood pressure (mmHg): intervention: SBP 138.9, (9.7); DBP 82.8, (7.7); control: SBP 139.9 (9.3); DBP 83.9, (5.8)</p>
Interventions	<p>Type: provision of foods high in fibre</p> <p>Description: Oat beta-glucans (intervention) group received a ready to eat cold cereal made with oat bran, oatmeal, and a powered form of oat B-glucans (soluble fibre).</p>

Maki 2007 (Continued)

	Control: a low-fibre, ready-to-eat, cold, wheat-based cereal, a low-fibre hot cereal and a control maltodextrin powder Both groups: participants instructed to eat three servings (one of each) per day Nutritional composition reported	
Outcomes	BP (no data reported)	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	High risk	High loss to follow-up
Selective reporting (reporting bias)	High risk	No data reported
Other bias	Unclear risk	Not enough information to judge

Makkonen 1993

Methods	Setting: Finland, setting not reported Design: individual randomisation, parallel group Dates: not reported Intervention duration: six months
Participants	N: 30 randomised (15 in each group). Baseline characteristics and results reported on different N's (where differ reported below) Inclusion criteria: women with climacteric symptoms, all normotensive. Hormonal therapy if used had been stopped for three months Exclusion criteria: no diabetes, liver disease or thromboembolic disease. Age (years): mean (SD); intervention: 52.3 (2.7); control: 53.6 (4.0) Sex (% men): 0% Ethnicity: not reported

	Cardiovascular risk status: mean (SD) BMI (kg/m²): not reported Total cholesterol (mmol/L): intervention: 7.51 (1.47), N = 15; control: 6.98 (0.63), N = 14 HDL cholesterol (mmol/L): intervention: 1.25 (0.39), N = 15; control: 1.31 (0.25), N = 12 LDL cholesterol (mmol/L): not reported Triglycerides (mmol/L): intervention: 1.74 (0.63), N = 15; control: 1.73 (0.83), N = 14 Blood pressure (mmHg): intervention: SBP 144.33 (21.66); DBP 92.6 (9.86) N = 15; control: SBP 148.8 (12.90); DBP 92.8 (7.89), N = 15	
Interventions	Type: fibre supplements Description: granulated guar gum (soluble fibre), 5 g three times per day. Control: placebo (wheat flour granules) 5 g three times per day	
Outcomes	BP, lipid levels, adverse events	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind and placebo
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No details on loss to follow-up or ITT
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	Unclear risk	Not enough information to judge

Marett 2004 Larch

Methods	Setting: USA, setting not reported Design: individual randomisation, parallel groups Dates: not reported Intervention duration: six months	
Participants	N: 54 randomised to three groups (intervention 1 N = 18; intervention 2, N = 19, control, N = 17) Inclusion criteria: healthy men and women, 18 years to 55 years Exclusion criteria: not stated Age (years): not reported Sex (% men): not reported Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): not reported Total cholesterol (mmol/L): intervention 1: 4.64 (1.00); intervention 2: 4.68 (0.99); control: 4.64 (1.04) HDL cholesterol (mmol/L): intervention 1: 1.25 (0.27); intervention 2: 1.32 (0.22); control: 1.34 (0.26) LDL cholesterol (mmol/L): intervention 1: 2.85 (1.00); intervention 2: 2.84 (0.96); control: 2.73 (0.83) Triglycerides (mmol/L): intervention 1: 1.17 (0.46); intervention 2: 1.02 (0.46); control: 1.21 (0.89) Blood pressure (mmHg): not reported	
Interventions	Type: fibre supplements Description: intervention 1: 8.4 g of larch arabinogalactan (soluble fibre); intervention 2: 8.4 g tamarack arabinogalactan (soluble fibre). control: 8.4 g placebo rice starch. Instructed to consume in a beverage or food	
Outcomes	BP, lipid levels	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported

Marett 2004 Larch (Continued)

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No details reported
Selective reporting (reporting bias)	High risk	Blood pressure is an outcome but no data
Other bias	Unclear risk	Not enough information to judge

Marett 2004 Tamarak

Methods	see previous - study had three relevant study groups	
Participants		
Interventions		
Outcomes		
Notes		

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	see previous
Allocation concealment (selection bias)	Unclear risk	see previous
Blinding of participants and personnel (performance bias) All outcomes	Low risk	see previous
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	see previous
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	see previous
Selective reporting (reporting bias)	High risk	see previous
Other bias	Unclear risk	see previous

Nichenametla 2014

Methods	<p>Setting: USA, community Design: cluster randomisation with a cross over Dates: not reported Intervention duration: 12 weeks</p>	
Participants	<p>N: 86 randomised (study results reported for total population only, numbers differ for baseline characteristics as reported below) Inclusion criteria: healthy participants and those with metabolic syndrome, 40.7% were taking medication for cholesterol, diabetes or blood pressure Exclusion criteria: pregnancy, lactation, long-term antibiotic therapy, immune compromised, cancer, other long term conditions that would affect the ability to provide informed consent or comply with protocol Age (years): mean (SD); all: 51.7 (14.8) Sex (% men): all: 31.4% Ethnicity: all: described as Caucasian 100% Cardiovascular risk status: mean (SD) BMI (kg/m²): all: 30.9 (5.1), N = 73 Total cholesterol (mmol/L): all: 5.2 (1.39), N = 83 HDL cholesterol (mmol/L): all: 1.4 (0.4), N = 83 LDL cholesterol (mmol/L): all: 3.4 (1.7), N = 75 Triglycerides (mmol/L): all: 1.2 (3.1), N = 83 Blood pressure (mmHg): all: SBP 131.6 (16.5), CBP 74.4 (10.1)</p>	
Interventions	<p>Type: provision of foods high in fibre Description: resistant starch-4 (RS4) an insoluble fibre, was substituted into the control flour. The intervention RS4-flour had a 30% v/v substitution of RS4 (Fibersym, MGP Ingredients, Atchison, KS) in the control flour. Participants consumed RS4-flour and control flour (without RS4 substitution) ad libitum in a free-living, domestic environment in form of any flour-based recipes that would normally be prepared, to match realistic conditions. Bread, noodles, maultaschen and dumplings are the most frequent flour based foods in Hutterite communities. Typical Hutterite diets contain high amounts of protein, fat, cholesterol and salt, but low fibre compared with RDA levels. A large number of food options are offered at each meal in a common dining hall. Nutrient composition (g/100 g) reported. Control flour. Details as above</p>	
Outcomes	BP, lipid levels, adverse events	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Two communities randomised as clusters
Allocation concealment (selection bias)	Unclear risk	No details reported

Nichenametla 2014 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind and treatments being blindly exchanged
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Unclear which groups drop outs relate to
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	High risk	Only two clusters randomised and evidence of a carry-over effect, although we only intended to use data from the first half of the trial before crossover

Pal 2011

Methods	<p>Setting: Australia, community setting</p> <p>Design: individual randomisation, parallel groups</p> <p>Dates: not reported</p> <p>Intervention duration: 12 weeks</p>
Participants	<p>N: 72 eligible participants (18 in each group). Baseline characteristics and results based on numbers after drop outs: fibre supplement: N = 16; control N = 15</p> <p>Inclusion criteria: overweight and obese, aged 18 years to 65 years</p> <p>Exclusion criteria: smoking, lipid lowering medication, steroids, agents that may influence lipid metabolism, warfarin, diabetes, hypo or hyperthyroidism, cardiovascular events in last six months, psychological unsuitability, major systemic disease, gastrointestinal problems, proteinuria, liver failure, renal failure, weight fluctuations six months, vegetarian</p> <p>Age (years): mean (SD); fibre supplement: 41.3 (9.2); control: 44.8 (6.2)</p> <p>Sex (% men): not reported</p> <p>Ethnicity: not reported</p> <p>Cardiovascular risk status: mean (SD)</p> <p>BMI (kg/m²): fibre supplement: 34.0 (3.6); control: 33.7 (3.9)</p> <p>Total cholesterol (mmol/L): fibre supplement: 5.9 (4.8); control: 6.1 (4.6)</p> <p>HDL cholesterol (mmol/L): fibre supplement: 1.3 (0.4); control: 1.3 (0.4)</p> <p>LDL cholesterol (mmol/L): fibre supplement: 4.0 (1.2); control: 3.9 (1.2)</p> <p>Triglycerides (mmol/L): fibre supplement: 1.5 (0.8); control: 2.0 (1.2)</p> <p>Blood pressure (mmHg): fibre supplement: SBP 115.0 (10.0); DBP 66.4 (10.4); control: SBP 114.3 (10.5); DBP 66.3 (8.9)</p>
Interventions	<p>Type: fibre supplement with usual diet</p> <p>Description: four groups: intervention 1: fibre supplement with usual diet, 12 g psyllium (soluble fibre); control: placebo with usual diet; Intervention 2: fibre supplement with</p>

	healthy eating advice, 12 g psyllium (not reported here); intervention 3: healthy eating advice with placebo (not reported here). Psyllium was mixed with 250 ml water and taken three times per day. The placebo consisted of breadcrumbs with flavouring to give 1.5 g soluble fibre	
Outcomes	BP, lipid levels	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	High risk	Single blinded study
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	High risk	High loss to follow up, no ITT, differences between groups
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	Unclear risk	Not enough information to judge

Perez-Jiminez 2008

Methods	<p>Setting: Spain, university setting</p> <p>Design: individual randomisation, parallel group</p> <p>Dates: not reported</p> <p>Intervention duration: 16 weeks</p>
Participants	<p>N: 43 randomised (intervention 34, control 9)</p> <p>Inclusion criteria: non-smokers, 25 were hypercholesteraemic (serum cholesterol > 200 mg/dl) and not taking any medication, mean age 33.7 (12.2) years</p> <p>Exclusion criteria: not reported</p> <p>Age (years): mean (SD); intervention: 35.5 (11.8); control: 34.6 (12.4)</p> <p>Sex (% men): intervention: 35.2%; control: 44.44%</p> <p>Ethnicity: not reported</p>

	<p>Cardiovascular risk status: mean (SD) BMI (kg/m²): intervention: 26.1 (4.7); control: 22.7 (2.4) Total cholesterol (mmol/L): intervention: 6.48 (1.67); control: 5.36 (1.13) HDL cholesterol (mmol/L): intervention: 1.23 (0.34); control: 1.1 (0.28) LDL cholesterol (mmol/L): intervention: 4.57 (1.40); control: 3.87 (0.9) Triglycerides (mmol/L): intervention: 1.36 (0.86); control: 0.85 (0.22) Blood pressure (mmHg): intervention: SBP 126.5 (22.1); DBP 78.2 (11.7) mmHg; control: SBP 121.5 (14.0); DBP 71.4 (14.4) mmHg</p>	
Interventions	<p>Type: fibre supplements Description: grape antioxidant dietary fibre (GADF) rich in dietary fibre and flavonoids (insoluble fibre), consumed for 16 weeks at 7.5 g daily in individual sterile packs, also consumed usual diet Control: consumed regular diet</p>	
Outcomes	BP, lipid levels, adverse events	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No details reported
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	High risk	Numbers reported without reasons, difference between groups
Selective reporting (reporting bias)	Unclear risk	No details reported
Other bias	Unclear risk	Not enough details to judge

Pins 2002

Methods	<p>Setting: USA, setting not reported Design: parallel randomised controlled trial Dates: not reported Intervention duration: 12 weeks</p>
Participants	<p>N: N = 88 randomised to two groups, 88 completed. Intervention group: N = 45, control group: N = 43 Inclusion criteria: mild or moderate hypertension treated with no more than one anti-hypertensive (excluding beta-blockers) Exclusion criteria: history of systolic BP > 160 mmHg or diastolic BP > 115 mmHg; existing complications of hypertension; history of major intestinal surgeries; malabsorption of the gastrointestinal tract or biliary disease; use of beta-blockers; diabetes mellitus; body mass index > 35; history of excessive use of alcohol; current smoking; high soluble fibre intake; clinical use of antacids, bulk laxatives or other medications affecting gastrointestinal tract; continuous treatment with oestrogen replacements, participation in another study three months before randomisation Age (years): control group 46.4 years (+/- 15.3 SD); intervention group 48.7 years (+/- 16.9 SD) Sex (% men): male/female ratio: 45/43 Ethnicity: 96%/98% (intervention/control) Caucasian Baseline cardiovascular risk status: <i>BMI (kg/m²):</i> intervention: 31.2 (5.1); control: 30.6 (4.7) <i>Total cholesterol (mmol/L):</i> intervention: 5.47 (0.99); control: 5.53 (1.09) <i>HDL cholesterol (mmol/L):</i> intervention: 1.11 (0.24); control: 1.14 (0.26) <i>LDL cholesterol (mmol/L):</i> intervention: 3.59 (0.76); control: 3.56 (0.71) <i>Triglycerides (mmol/L):</i> intervention: 2.09 (0.45); control: 2.16 (0.47) Blood pressure (mmHg): intervention: SBP 140 (16); DBP 88 (10) mmHg; control: SBP 138 (15); DBP 86 (9) mmHg Medications used: no more than one anti-hypertensive medication and/or one diuretic medication. Eighty participants were on a single anti-hypertensive medication, eight were on an anti-hypertensive drug plus a diuretic medication</p>
Interventions	<p>Type: provision of foods high in fibre Description: whole oat cereals (Quaker Oatmeal and Oat Squares) eaten as part of usual diet The control group received refined grain wheat-based cereals (hot wheat cereal and Kellogg's Crispix) eaten as part of usual diet. The intervention group received the following: 60 g Quaker oatmeal (5.61 g total dietary fibre, 3.25 g soluble fibre and 2.83 g beta-glucans) and 77 g Quaker Oatmeal Squares (6.07 g total fibre, 2.98 g soluble fibre, 2.59 g beta-glucans). The control group received the following: 65 g Malt-O-Meal hot wheat cereal (2.32 g total dietary fibre, 0.6 g soluble fibre) and 81 g Kelloggs Crispix (1.2 g dietary fibre, < 0.5 g soluble fibre). Cereal treatments were isocaloric</p>
Outcomes	<p>Total cholesterol, HDL cholesterol, LDL cholesterol, triglycerides. Proportion of participants reducing anti-hypertensive medication (no variance provided), mean BP (no variance provided). Adverse events</p>
Notes	<p>Test cereals and funding were provided by the Quaker Oats Company All participants were on anti-hypertensive treatment at the start of the study. This was reduced by half at four weeks and fully at 12 weeks if participants' blood pressure was</p>

Pins 2002 (Continued)

	within certain limits. We have not used the blood pressure data from this trial because of the varying effects of antihypertensive medication throughout the trial confounding any effects of fibre	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not reported
Allocation concealment (selection bias)	Unclear risk	Not reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Cereals were dispensed in unlabelled bulk containers to facilitate blinding
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The outcome assessor was blinded to group allocation
Incomplete outcome data (attrition bias) All outcomes	Low risk	All 88 subjects randomised appear to have completed the study
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	High risk	Study was funded by the Quaker Oats Company who also provided cereals for the intervention

Reimer 2013

Methods	<p>Setting: Japan, setting not reported</p> <p>Design: individual randomisation, parallel group</p> <p>Dates: not reported</p> <p>Intervention duration: 14 weeks</p>
Participants	<p>N: 64 randomised (N = 32 to each group). Baseline characteristics and results based on N = 28 for both groups</p> <p>Inclusion criteria: BMI 24-30 kg/m² and waist circumference > 85 cm (males) or > 90 cm (females), aged 20 years to 65 years</p> <p>Exclusion criteria: heavier than 100 kg, attempting to lose weight, taking medications that alter blood lipids or blood glucose, under medically supervised treatments for hyperlipidaemia, diabetes or obesity, history of allergy to test products, disease that requires medication, participation in another trial, pregnancy, breast feeding</p> <p>Age (years): not reported</p> <p>Sex (% men): 43.7% for both groups (randomised N = 64)</p>

	<p>Ethnicity: not reported</p> <p>Cardiovascular risk status: mean (SD)</p> <p>BMI (kg/m²): intervention: 26.7 (1.1); control: 27.2 (1.6)</p> <p>Total cholesterol (mmol/L): intervention: 5.63 (1.1); control: 5.32 (0.8)</p> <p>HDL cholesterol (mmol/L): intervention: 1.45 (0.3); control: 1.39 (0.3)</p> <p>LDL cholesterol (mmol/L): intervention: 3.69 (0.8); control: 3.41 (0.7)</p> <p>Triglycerides (mmol/L): intervention: 1.38 (0.8); control: 1.29 (0.7)</p> <p>Blood pressure (mmHg): intervention: SBP 123.6 (11.1); DBP 75.1 (6.9); control: SBP 120.2 (14.3); DBP 75.1 (13.2)</p>	
Interventions	<p>Type: fibre supplements</p> <p>Description: PolyGlycoplex (PGX) (soluble fibre). Provided in packets labelled A or B, five minutes before each meal, 5 g of PGX was mixed with yogurt. Days one to three, 5 g per day, days four to seven, 10 g per day, days eight to 97, 15 g per day. Control: same as intervention but with rice flour placebo</p>	
Outcomes	BP, lipid levels, adverse events	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Used placebo, states study coordinator and participants blinded to product
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Loss to follow-up reported with reasons
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	Unclear risk	Not enough information to judge

Schlamovitz 1987

Methods	Setting: Denmark, setting not reported Design: individual randomisation, parallel group Dates: not reported Intervention duration: 12 weeks	
Participants	N: 46 randomised (Intervention 21, Control 25) Inclusion criteria: mild to moderate hypertension (BP between 140/95 and 198/110). Described as 'lean' patients. No further details Exclusion criteria: not reported Age (years): not reported Sex (% men): not reported Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): not reported Total cholesterol (mmol/L): not reported HDL cholesterol (mmol/L): intervention: 5.9 (1.3); control: 6.5 (1.2) LDL cholesterol (mmol/L): not reported Triglycerides (mmol/L): not reported Blood pressure (mmHg): intervention: SBP 157 (17.6), DBP 97 (6.6); control: SBP 150 (12.1), DBP 96 (7.3)	
Interventions	Type: fibre supplements Description: FibreTrim Plus tablets, Farma Food (soluble fibre). Tablets contain about 58% fibre from grain, citrus fruit, and vegetables. Total dietary fibre supplement 7 g per day. Control given placebo (no further details)	
Outcomes	BP, lipid levels, adverse events	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No details reported
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind, objective outcome measures
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No details reported

Selective reporting (reporting bias)	High risk	Letter with limited information, however, triglycerides data missing
Other bias	Unclear risk	Not enough information to judge

Shimizu 2008

Methods	Setting: Japan, setting not stated Design: individual randomisation, parallel group Dates: not reported Intervention duration: 12 weeks	
Participants	N: 44 randomised (22 to each group). Baseline characteristics and results based on numbers after drop out (intervention N = 19, control N = 20) Inclusion criteria: BMI > 22 Kg/m ² , LDL-C 140-220 mg/dl. Total C 220-300 mg/dl. Habitually consumed pearled rice as a staple food more than twice in three daily meals, age 30 years to 60 years Exclusion criteria: Heart, liver, kidney disease, food allergies, exercise habits daily, body weight increase or decrease by > 10 kg in past three months, irregular lifestyle habits, medication or functional foods known to affect lipid metabolism Age (years): mean (SD); intervention: 42.1 (9.2) control: 40.9 (7.8) Sex (% men): not reported Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): intervention: 26.2 (2.8); control: 24.5 (2.4) Total cholesterol (mmol/L): intervention: 6.07 (0.56); control: 6.36 (0.70) HDL cholesterol (mmol/L): not reported LDL cholesterol (mmol/L): intervention: 3.97 (0.42); control: 4.14 (0.62) Triglycerides (mmol/L) not reported Blood pressure (mmHg): not reported	
Interventions	Type: provision of foods high in fibre Description: High beta-glucagon barley (insoluble fibre), each package contained 50% barley and 50% rice in a package (160 g) and participants consumed 2 packages daily in replacement of the pearled rice in their staple food. The dietary fibre content of each package was 4.5 g of which beta-glucagon was 3.5 g per package (so 7g per day in total). The control group consumed rice only in a package (139 g) and consumed two packages daily	
Outcomes	Lipid levels	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement

Shimizu 2008 (Continued)

Random sequence generation (selection bias)	Unclear risk	States randomised on basis of baseline factors but no other details
Allocation concealment (selection bias)	Unclear risk	No details reported
Blinding of participants and personnel (performance bias) All outcomes	Low risk	States double blind, same packaging used
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	No details reported
Incomplete outcome data (attrition bias) All outcomes	High risk	Dropouts not described per study group
Selective reporting (reporting bias)	Unclear risk	All outcomes stated were reported but unable to fully judge
Other bias	Unclear risk	Not enough information to judge

Tighe 2010 Wheat

Methods	<p>Setting: UK, setting not described Design: individual randomisation, parallel group Dates: June 2005 to September 2008 Intervention duration: 12 weeks</p>
Participants	<p>N: 226 randomised (intervention 1: N = 77; intervention 2: N = 73; control: N = 76). Baseline characteristics and results based on Intervention 1: N = 73; intervention 2: N = 70; control: N = 63 Inclusion criteria: BMI 18.5-35 kg/m², sedentary or moderately active (fewer than two aerobic sessions per week), aged 40-60 years, signs of metabolic syndrome or moderate hypercholesterolaemia Exclusion criteria: CVD, diabetes, fasting blood glucose > 7 mmol/L, asthma, SBP > 160 mmHg or DBP > 99 mmHg, thyroid or eating disorders, high habitual intake of whole grain foods, regular medication or supplements known to affect any outcomes Age (years): mean (SD); intervention 1:51.6 (7.4); intervention 2:52.1 (7.4); control: 51.8 (7.4) Sex (% men): intervention 1: 53%; intervention 2: 51%; control: 47% Ethnicity: not reported Cardiovascular risk status: mean (SD) BMI (kg/m²): intervention 1: 28.0 (4.2); intervention 2: 27.0 (3.7); control: 28.0 (4.0) Total cholesterol (mmol/L): intervention 1: 5.46 (1.18); intervention 2: 5.57 (1.03); control: 5.94 (1.11) HDL cholesterol (mmol/L): intervention 1: 1.55 (0.40); intervention 2: 1.62 (0.40); control: 1.62 (0.48) LDL cholesterol (mmol/L): intervention 1: 3.45 (1.03); intervention 2: 3.45 (0.89);</p>

Tighe 2010 Wheat (Continued)

	control: 3.66 (0.98) <i>Triglycerides (mmol/L)</i> : intervention 1: 1.27 (0.68); intervention 2: 1.12 (0.54); control: 1.49 (0.86) <i>Blood pressure (mmHg)</i> : intervention 1: SBP 125.9 (1.4); DBP 75.7 (0.8); intervention 2: SBP 131.7 (1.4); DBP 78.4 (0.8); control: SBP 131.2 (1.4); DBP 79.1 (0.8)	
Interventions	Type: provision of foods high in fibre Description: intervention 1: three servings of whole wheat foods (70 g to 80 g wholemeal bread and 30 g to 40 g whole grain cereals) (insoluble fibre). Intervention 2: one serving of whole wheat foods (35 g to 40 g wholemeal bread) and two servings of oats (60 g to 80 g of whole grain rolled oats) (soluble and insoluble fibre). Control: three servings of refined cereal foods (refined cereals and white bread). All instructed not to alter food intake apart from the trial changes and to maintain usual level of activity	
Outcomes	BP, lipid levels	
Notes	With thanks also to Dr Frank Theis for providing additional data for their trial (Tighe 2010 Wheat ; Tighe 2010 Wheat+Oats).	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Web-based randomisation system, random permuted blocks stratified by age, gender and BMI
Allocation concealment (selection bias)	Low risk	Off-site allocation
Blinding of participants and personnel (performance bias) All outcomes	High risk	Single blind
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome assessors blind
Incomplete outcome data (attrition bias) All outcomes	High risk	Numbers and reasons provided, difference between groups
Selective reporting (reporting bias)	Unclear risk	Blood pressure reported in Tighe 2010 paper, not 2013 paper
Other bias	Unclear risk	Not enough information to judge

Tighe 2010 Wheat+Oats

Methods	see previous - study had three relevant study groups	
Participants		
Interventions		
Outcomes		
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	see previous
Allocation concealment (selection bias)	Low risk	see previous
Blinding of participants and personnel (performance bias) All outcomes	High risk	see previous
Blinding of outcome assessment (detection bias) All outcomes	Low risk	see previous
Incomplete outcome data (attrition bias) All outcomes	High risk	see previous
Selective reporting (reporting bias)	Unclear risk	see previous
Other bias	Unclear risk	see previous

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Bays 2013	Short term (six weeks)
Davy 2002	Control not no intervention or minimal intervention
Dodson 1989	Multifactorial intervention
King 2008	No relevant outcomes

(Continued)

Liao 2007	Short term (eight weeks)
Mee 1997	Short term (six weeks)
Ramprasath 2014	Multifactorial intervention
Reid 2002	Control not no intervention or minimal intervention
Tapola 2008	Short term (eight weeks)

Characteristics of studies awaiting assessment [ordered by study ID]

Keenan 2002b

Methods	Cost analysis of a parallel group RCT included in the review (Pins 2002)
Participants	see Pins 2002
Interventions	see Pins 2002
Outcomes	see Pins 2002
Library unable to locate	Yes
Notes	This is the abstract from the database search: The management of hypertension has become an increasing priority in managed care. While better control of blood pressure will lead to better health and, we expect, lower costs in the future, the short term cost of antihypertensive medication are here and now. In this report, Joseph Keenan and his colleagues demonstrate lower costs for antihypertensive medication in patients randomly assigned to a diet rich in whole-grain fiber, compared to controls. While it seems appropriate to call for further study and longer follow-up, these findings could form the basis for much more cost-effective approaches to managing the hypertensive population

Characteristics of ongoing studies [ordered by study ID]

Lambert 2014

Trial name or title	
Methods	Parallel group RCT
Participants	Stable body weight for at least three months, aged 18 years to 70 years, BMI between 25 kg/m ² to 58 kg/m ² , no concomitant weight loss medication use, diet or exercise regime for weight loss, corticosteroids, antidepressants, anti-epileptics, lipid lowering or diabetes medications, previous bariatric surgery or other intestinal surgery, pregnancy or lactation, use of bulk laxatives, probiotics, chronic antacid use, antibiotics, significant cardiovascular or respiratory disease, liver disease, alcohol or drug dependency, active malignancy,

Lambert 2014 (Continued)

	> 350 lb weight
Interventions	Twelve week intervention of food high in fibre Biscuits containing 5 g of pea fibre per serving, three times per day within 30 minutes of daily meals. Dose slowly increased over the first three weeks of the study Control group to have an isocaloric control biscuit with no pea fibre
Outcomes	Lipid levels
Starting date	October 2012
Contact information	reimer@ucalgary.ca
Notes	ClinicalTrials.gov (NCT01719900) Registered October 23, 2012

DATA AND ANALYSES

Comparison 1. Fibre versus control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Total Cholesterol mmol/L change	20	1067	Mean Difference (IV, Random, 95% CI)	-0.23 [-0.40, -0.06]
2 HDL Cholesterol mmol/L change	18	982	Mean Difference (IV, Fixed, 95% CI)	-0.03 [-0.06, -0.01]
3 LDL Cholesterol mmol/L change	18	995	Mean Difference (IV, Fixed, 95% CI)	-0.14 [-0.22, -0.06]
4 Triglycerides mmol/L change	18	982	Mean Difference (IV, Fixed, 95% CI)	0.00 [-0.04, 0.05]
5 Systolic blood pressure (mmHg) change	10	661	Mean Difference (IV, Random, 95% CI)	-1.92 [-4.02, 0.19]
6 Diastolic blood pressure (mmHg) change	10	661	Mean Difference (IV, Fixed, 95% CI)	-1.77 [-2.61, -0.92]

Comparison 2. Subgroup analyses

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Total cholesterol mmol/L change	20	1067	Mean Difference (IV, Random, 95% CI)	-0.23 [-0.40, -0.06]
1.1 Type of intervention - fibre supplement	13	555	Mean Difference (IV, Random, 95% CI)	-0.27 [-0.50, -0.04]
1.2 Type of Intervention - provision of foods	7	512	Mean Difference (IV, Random, 95% CI)	-0.16 [-0.42, 0.09]
2 HDL Cholesterol mmol/L change	18	982	Mean Difference (IV, Fixed, 95% CI)	-0.03 [-0.06, -0.01]
2.1 Type of intervention - fibre supplement	12	509	Mean Difference (IV, Fixed, 95% CI)	-0.04 [-0.07, -0.00]
2.2 Type of intervention - provision of foods	6	473	Mean Difference (IV, Fixed, 95% CI)	-0.03 [-0.07, 0.02]
3 LDL Cholesterol mmol/L change	18	995	Mean Difference (IV, Fixed, 95% CI)	-0.14 [-0.22, -0.06]
3.1 Type of Intervention - fibre supplement	11	483	Mean Difference (IV, Fixed, 95% CI)	-0.13 [-0.24, -0.03]
3.2 Type of Intervention - provision of foods	7	512	Mean Difference (IV, Fixed, 95% CI)	-0.14 [-0.26, -0.03]
4 Triglycerides mmol/L change	18	982	Mean Difference (IV, Fixed, 95% CI)	0.00 [-0.04, 0.05]
4.1 Type of Intervention - fibre supplement	12	509	Mean Difference (IV, Fixed, 95% CI)	-0.03 [-0.12, 0.05]
4.2 Type of Intervention - provision of foods	6	473	Mean Difference (IV, Fixed, 95% CI)	0.02 [-0.03, 0.06]


5	Systolic blood pressure (mmHg) change	10	661	Mean Difference (IV, Random, 95% CI)	-1.92 [-4.02, 0.19]
5.1	Type of intervention - fibre supplement	6	310	Mean Difference (IV, Random, 95% CI)	-2.14 [-5.19, 0.91]
5.2	Type of intervention - provision of foods	4	351	Mean Difference (IV, Random, 95% CI)	-1.57 [-4.45, 1.31]
6	Diastolic blood pressure (mmHg) change	10	653	Mean Difference (IV, Fixed, 95% CI)	-1.77 [-2.62, -0.92]
6.1	Type of intervention - fibre supplement	6	310	Mean Difference (IV, Fixed, 95% CI)	-1.07 [-2.30, 0.16]
6.2	Type of intervention - provision of foods	4	343	Mean Difference (IV, Fixed, 95% CI)	-2.40 [-3.57, -1.23]
7	Total cholesterol mmol/L change	20	1067	Mean Difference (IV, Random, 95% CI)	-0.23 [-0.40, -0.06]
7.1	Type of fibre - soluble	14	688	Mean Difference (IV, Random, 95% CI)	-0.26 [-0.46, -0.05]
7.2	Type of fibre - insoluble	3	187	Mean Difference (IV, Random, 95% CI)	-0.03 [-0.53, 0.47]
7.3	Type of fibre - soluble and insoluble	3	192	Mean Difference (IV, Random, 95% CI)	-0.28 [-0.75, 0.19]
8	HDL Cholesterol mmol/L change	18	982	Mean Difference (IV, Fixed, 95% CI)	-0.03 [-0.06, -0.01]
8.1	Type of fibre - soluble	13	642	Mean Difference (IV, Fixed, 95% CI)	-0.03 [-0.06, 0.00]
8.2	Type of fibre - insoluble	2	148	Mean Difference (IV, Fixed, 95% CI)	0.00 [-0.13, 0.14]
8.3	Type of fibre - soluble and insoluble	3	192	Mean Difference (IV, Fixed, 95% CI)	-0.07 [-0.14, 0.00]
9	LDL Cholesterol mmol/L change	18	995	Mean Difference (IV, Fixed, 95% CI)	-0.14 [-0.22, -0.06]
9.1	Type of fibre - soluble	12	616	Mean Difference (IV, Fixed, 95% CI)	-0.14 [-0.23, -0.05]
9.2	Type of fibre - insoluble	3	187	Mean Difference (IV, Fixed, 95% CI)	0.01 [-0.24, 0.27]
9.3	Type of fibre - soluble and insoluble	3	192	Mean Difference (IV, Fixed, 95% CI)	-0.25 [-0.48, -0.02]
10	Triglycerides mmol/L change	18	982	Mean Difference (IV, Fixed, 95% CI)	0.00 [-0.04, 0.05]
10.1	Type of fibre - soluble	13	642	Mean Difference (IV, Fixed, 95% CI)	-0.05 [-0.13, 0.03]
10.2	Type of fibre - insoluble	2	148	Mean Difference (IV, Fixed, 95% CI)	-0.11 [-0.32, 0.10]
10.3	Type of fibre - soluble and insoluble	3	192	Mean Difference (IV, Fixed, 95% CI)	0.03 [-0.02, 0.08]
11	Systolic blood pressure (mmHg) change	10	661	Mean Difference (IV, Random, 95% CI)	-1.92 [-4.02, 0.19]
11.1	Type of fibre - soluble	6	377	Mean Difference (IV, Random, 95% CI)	-2.19 [-4.66, 0.28]
11.2	Type of fibre -insoluble	2	146	Mean Difference (IV, Random, 95% CI)	-3.19 [-7.91, 1.52]
11.3	Type of fibre - soluble and insoluble	2	138	Mean Difference (IV, Random, 95% CI)	-1.26 [-7.50, 4.98]
12	Diastolic blood pressure (mmHg) change	10	661	Mean Difference (IV, Fixed, 95% CI)	-1.77 [-2.61, -0.92]
12.1	Type of fibre - soluble	6	377	Mean Difference (IV, Fixed, 95% CI)	-1.04 [-2.15, 0.07]
12.2	Type of fibre -insoluble	2	146	Mean Difference (IV, Fixed, 95% CI)	-1.09 [-3.97, 1.79]
12.3	Type of fibre - soluble and insoluble	2	138	Mean Difference (IV, Fixed, 95% CI)	-3.21 [-4.67, -1.74]

Analysis 1.1. Comparison 1 Fibre versus control, Outcome 1 Total Cholesterol mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease

Comparison: 1 Fibre versus control

Outcome: 1 Total Cholesterol mmol/L change




Analysis 1.2. Comparison 1 Fibre versus control, Outcome 2 HDL Cholesterol mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease

Comparison: 1 Fibre versus control

Outcome: 2 HDL Cholesterol mmol/L change




Analysis 1.3. Comparison 1 Fibre versus control, Outcome 3 LDL Cholesterol mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease

Comparison: 1 Fibre versus control

Outcome: 3 LDL Cholesterol mmol/L change




Analysis 1.4. Comparison 1 Fibre versus control, Outcome 4 Triglycerides mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease

Comparison: 1 Fibre versus control

Outcome: 4 Triglycerides mmol/L change




Analysis 1.5. Comparison 1 Fibre versus control, Outcome 5 Systolic blood pressure (mmHg) change.

Review: Dietary fibre for the primary prevention of cardiovascular disease

Comparison: 1 Fibre versus control

Outcome: 5 Systolic blood pressure (mmHg) change




Analysis 1.6. Comparison 1 Fibre versus control, Outcome 6 Diastolic blood pressure (mmHg) change.

Review: Dietary fibre for the primary prevention of cardiovascular disease

Comparison: 1 Fibre versus control

Outcome: 6 Diastolic blood pressure (mmHg) change




Analysis 2.1. Comparison 2 Subgroup analyses, Outcome 1 Total cholesterol mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease


Comparison: 2 Subgroup analyses

Outcome: 1 Total cholesterol mmol/L change



(Continued ...)

(... Continued)




Analysis 2.2. Comparison 2 Subgroup analyses, Outcome 2 HDL Cholesterol mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease


Comparison: 2 Subgroup analyses

Outcome: 2 HDL Cholesterol mmol/L change



(Continued ...)

(... Continued)




Analysis 2.3. Comparison 2 Subgroup analyses, Outcome 3 LDL Cholesterol mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease

Comparison: 2 Subgroup analyses

Outcome: 3 LDL Cholesterol mmol/L change




Analysis 2.4. Comparison 2 Subgroup analyses, Outcome 4 Triglycerides mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease


Comparison: 2 Subgroup analyses

Outcome: 4 Triglycerides mmol/L change



(Continued ...)

(... Continued)




Analysis 2.5. Comparison 2 Subgroup analyses, Outcome 5 Systolic blood pressure (mmHg) change.

Review: Dietary fibre for the primary prevention of cardiovascular disease


Comparison: 2 Subgroup analyses

Outcome: 5 Systolic blood pressure (mmHg) change



(Continued ...)

(... Continued)




Analysis 2.6. Comparison 2 Subgroup analyses, Outcome 6 Diastolic blood pressure (mmHg) change.

Review: Dietary fibre for the primary prevention of cardiovascular disease

Comparison: 2 Subgroup analyses

Outcome: 6 Diastolic blood pressure (mmHg) change




Analysis 2.7. Comparison 2 Subgroup analyses, Outcome 7 Total cholesterol mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease


Comparison: 2 Subgroup analyses

Outcome: 7 Total cholesterol mmol/L change



(Continued ...)

(... Continued)




Analysis 2.8. Comparison 2 Subgroup analyses, Outcome 8 HDL Cholesterol mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease


Comparison: 2 Subgroup analyses

Outcome: 8 HDL Cholesterol mmol/L change



(Continued ...)

(... Continued)




Analysis 2.9. Comparison 2 Subgroup analyses, Outcome 9 LDL Cholesterol mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease


Comparison: 2 Subgroup analyses

Outcome: 9 LDL Cholesterol mmol/L change



(Continued ...)

(... Continued)




Analysis 2.10. Comparison 2 Subgroup analyses, Outcome 10 Triglycerides mmol/L change.

Review: Dietary fibre for the primary prevention of cardiovascular disease


Comparison: 2 Subgroup analyses

Outcome: 10 Triglycerides mmol/L change



(Continued ...)

(... Continued)




Analysis 2.11. Comparison 2 Subgroup analyses, Outcome 11 Systolic blood pressure (mmHg) change.

Review: Dietary fibre for the primary prevention of cardiovascular disease

Comparison: 2 Subgroup analyses

Outcome: 11 Systolic blood pressure (mmHg) change




Analysis 2.12. Comparison 2 Subgroup analyses, Outcome 12 Diastolic blood pressure (mmHg) change.

Review: Dietary fibre for the primary prevention of cardiovascular disease

Comparison: 2 Subgroup analyses

Outcome: 12 Diastolic blood pressure (mmHg) change



APPENDICES

Appendix I. Search Strategy

Cochrane Library

#1MeSH descriptor: [Dietary Fiber] explode all trees

#2roughage*

#3prebiotic*

#4carbohydrate near/2 polymer*

#5((non-starch or nonstarch) near (poly-saccharide* or polysaccharide*))

#6#1 or #2 or #3 or #4 or #5

#7MeSH descriptor: [Diet] this term only

#8diet*

#9consum*

#10eat*

#11food*

#12nutri*

#13#7 or #8 or #9 or #10 or #11 or #12

#14MeSH descriptor: [Chitin] explode all trees

#15MeSH descriptor: [Cellulose] explode all trees

#16chitin*

#17lignin*

#18hemicellulose*

#19hexosane*

#20pentosane*

#21xanthan*

#22MeSH descriptor: [Fructans] explode all trees

#23fructan*

#24inulin*

#25polyuronide*

#26MeSH descriptor: [Pectins] explode all trees

#27pectin*

#28MeSH descriptor: [Alginates] this term only

#29alginate*

#30alginic near/2 acid*

#31MeSH descriptor: [Agar] this term only

#32agar*

#33MeSH descriptor: [Carrageenan] this term only

#34carrageen*

#35MeSH descriptor: [Raffinose] this term only

#36raffinose*

#37MeSH descriptor: [Xylose] this term only

#38xylose*

#39polydextrose*

#40MeSH descriptor: [Lactulose] this term only

#41lactulose*

#42cellulose*

#43fibre* or fiber* or high-fibre* or high-fiber*

#44#14 or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28 or #29 or #30 or #31 or #32 or #33 or #34 or #35 or #36 or #37 or #38 or #39 or #40 or #41 or #42 or #43

#45#13 and #44

#46#6 or #45
 #47MeSH descriptor: [Cardiovascular Diseases] explode all trees
 #48cardio*
 #49cardia*
 #50heart*
 #51coronary*
 #52angina*
 #53ventric*
 #54myocard*
 #55pericard*
 #56isch?em*
 #57emboli*
 #58arrhythmi*
 #59thrombo*
 #60atrial next fibrillat*
 #61tachycardi*
 #62endocardi*
 #63(sick next sinus)
 #64MeSH descriptor: [Stroke] explode all trees
 #65(stroke or stokes)
 #66cerebrovasc*
 #67cerebral next vascular
 #68apoplexy
 #69(brain near/2 accident*)
 #70((brain* or cerebral or lacunar) near/2 infarct*)
 #71MeSH descriptor: [Hypertension] explode all trees
 #72hypertensi*
 #73(peripheral next arter* next disease*)
 #74((high or increased or elevated) near/2 blood pressure)
 #75MeSH descriptor: [Hyperlipidemias] explode all trees
 #76hyperlipid*
 #77hyperlip?emia*
 #78hypercholesterol*
 #79hypercholester?emia*
 #80hyperlipoprotein?emia*
 #81hypertriglycerid?emia*
 #82MeSH descriptor: [Arteriosclerosis] explode all trees
 #83MeSH descriptor: [Cholesterol] explode all trees
 #84cholesterol
 #85“coronary risk factor*”
 #86MeSH descriptor: [Blood Pressure] this term only
 #87“blood pressure”
 #88#47 or #48 or #49 or #50 or #51 or #52 or #53 or #54 or #55 or #56 or #57 or #58 or #59 or #60 or #61 or #62 or #63 or #64 or #65 or #66 or #67 or #68 or #69 or #70 or #71 or #72 or #73 or #74 or #75 or #76 or #77 or #78 or #79 or #80 or #81 or #82 or #83 or #84 or #85 or #86 or #87
 #89#46 and #88

MEDLINE OVID

1. exp Dietary Fiber/
2. roughage*.tw.
3. prebiotic*.tw.
4. (carbohydrate adj2 polymer*).tw.

5. ((non-starch or nonstarch) adj (poly-saccharide* or polysaccharide*)).tw.
6. or/1-5
7. Diet/
8. diet*.tw.
9. consum*.tw.
10. eat*.tw.
11. food*.tw.
12. nutri*.tw.
13. or/7-12
14. exp Chitin/
15. exp Cellulose/
16. chitin*.tw.
17. lignin*.tw.
18. hemicellulose*.tw.
19. hexosane*.tw.
20. pentosane*.tw.
21. xanthan*.tw.
22. exp Fructans/
23. fructan*.tw.
24. inulin*.tw.
25. polyuronide*.tw.
26. Pectins/
27. pectin*.tw.
28. Alginates/
29. alginate*.tw.
30. (alginic adj2 acid*).tw.
31. Agar/
32. agar*.tw.
33. Carrageenan/
34. carrageen*.tw.
35. Raffinose/
36. raffinose*.tw.
37. Xylose/
38. xylose*.tw.
39. polydextrose*.tw.
40. Lactulose/
41. lactulose*.tw.
42. cellulose*.tw.
43. (fibre* or fiber* or high-fibre* or high-fiber*).tw.
44. or/14-43
45. 13 and 44
46. 6 or 45
47. exp Cardiovascular Diseases/
48. cardio*.tw.
49. cardia*.tw.
50. heart*.tw.
51. coronary*.tw.
52. angina*.tw.
53. ventric*.tw.
54. myocard*.tw.
55. pericard*.tw.
56. isch?em*.tw.
57. emboli*.tw.

58. arrhythmi*.tw.
59. thrombo*.tw.
60. atrial fibrillat*.tw.
61. tachycardi*.tw.
62. endocardi*.tw.
63. (sick adj sinus).tw.
64. exp Stroke/
65. (stroke or stokes).tw.
66. cerebrovasc*.tw.
67. cerebral vascular.tw.
68. apoplexy.tw.
69. (brain adj2 accident*).tw.
70. ((brain* or cerebral or lacunar) adj2 infarct*).tw.
71. exp Hypertension/
72. hypertensi*.tw.
73. peripheral arter* disease*.tw.
74. ((high or increased or elevated) adj2 blood pressure).tw.
75. exp Hyperlipidemias/
76. hyperlipid*.tw.
77. hyperlip?emia*.tw.
78. hypercholesterol*.tw.
79. hypercholester?emia*.tw.
80. hyperlipoprotein?emia*.tw.
81. hypertriglycerid?emia*.tw.
82. exp Arteriosclerosis/
83. exp Cholesterol/
84. cholesterol.tw.
85. "coronary risk factor* ".tw.
86. Blood Pressure/
87. blood pressure.tw.
88. or/47-87
89. 46 and 88
90. randomized controlled trial.pt.
91. controlled clinical trial.pt.
92. randomized.ab.
93. placebo.ab.
94. drug therapy.fs.
95. randomly.ab.
96. trial.ab.
97. groups.ab.
98. 90 or 91 or 92 or 93 or 94 or 95 or 96 or 97
99. exp animals/ not humans.sh.
100. 98 not 99
101. 89 and 100

EMBASE OVID

1. dietary fiber/
2. prebiotic agent/
3. roughage*.tw.
4. prebiotic*.tw.
5. (carbohydrate adj2 polymer*).tw.
6. ((non-starch or nonstarch) adj (poly-saccharide* or polysaccharide*)).tw.

7. high fiber diet/
8. or/1-7
9. diet/
10. diet*.tw.
11. consum*.tw.
12. eat*.tw.
13. food*.tw.
14. nutri*.tw.
15. or/9-14
16. chitin/
17. chitosan/
18. cellulose/
19. lignin/
20. chitin*.tw.
21. lignin*.tw.
22. hemicellulose*.tw.
23. hexosane*.tw.
24. pentosane*.tw.
25. xanthan*.tw.
26. xanthan/
27. fructan/
28. inulin/
29. fructan*.tw.
30. inulin*.tw.
31. polyuronide*.tw.
32. pectin/
33. pectin*.tw.
34. alginic acid/
35. alginate*.tw.
36. (alginic adj2 acid*).tw.
37. agar/
38. agar*.tw.
39. carrageenan/
40. carrageen*.tw.
41. raffinose/
42. raffinose*.tw.
43. xylose/
44. xylose*.tw.
45. polydextrose*.tw.
46. lactulose/
47. lactulose*.tw.
48. cellulose*.tw.
49. (fibre* or fiber* or high-fibre* or high-fiber*).tw.
50. or/16-49
51. 15 and 50
52. 8 or 51
53. exp cardiovascular disease/
54. cardio*.tw.
55. cardia*.tw.
56. heart*.tw.
57. coronary*.tw.
58. angina*.tw.
59. ventric*.tw.

60. myocard*.tw.
61. pericard*.tw.
62. isch?em*.tw.
63. emboli*.tw.
64. arrhythmi*.tw.
65. thrombo*.tw.
66. atrial fibrillat*.tw.
67. tachycardi*.tw.
68. endocardi*.tw.
69. (sick adj sinus).tw.
70. exp cerebrovascular disease/
71. (stroke or stokes).tw.
72. cerebrovasc*.tw.
73. cerebral vascular.tw.
74. apoplexy.tw.
75. (brain adj2 accident*).tw.
76. ((brain* or cerebral or lacunar) adj2 infarct*).tw.
77. exp hypertension/
78. hypertensi*.tw.
79. peripheral arter* disease*.tw.
80. ((high or increased or elevated) adj2 blood pressure).tw.
81. exp hyperlipidemia/
82. hyperlipid*.tw.
83. hyperlip?emia*.tw.
84. hypercholesterol*.tw.
85. hypercholester?emia*.tw.
86. hyperlipoprotein?emia*.tw.
87. hypertriglycerid?emia*.tw.
88. exp Arteriosclerosis/
89. exp Cholesterol/
90. cholesterol.tw.
91. "coronary risk factor*".tw.
92. Blood Pressure/
93. blood pressure.tw.
94. or/53-93
95. 52 and 94
96. random\$.tw.
97. factorial\$.tw.
98. crossover\$.tw.
99. cross over\$.tw.
100. cross-over\$.tw.
101. placebo\$.tw.
102. (doubl\$ adj blind\$).tw.
103. (singl\$ adj blind\$).tw.
104. assign\$.tw.
105. allocat\$.tw.
106. volunteer\$.tw.
107. crossover procedure/
108. double blind procedure/
109. randomized controlled trial/
110. single blind procedure/
111. 96 or 97 or 98 or 99 or 100 or 101 or 102 or 103 or 104 or 105 or 106 or 107 or 108 or 109 or 110
112. (animal/ or nonhuman/) not human/

113. 111 not 112
 114. 95 and 113
 115. limit 114 to embase

Web of Science

- # 16 #15 AND #14
 # 15 TS=(random* or blind* or allocat* or assign* or trial* or placebo* or crossover* or cross-over*)
 # 14 #13 AND #12
 # 13 #10 OR #9 OR #8 OR #7 OR #6 OR #5 OR #4
 # 12 #11 OR #1
 # 11 #3 AND #2
 # 10 TS=(hyperlipid* OR hyperlip?emia* OR hypercholesterol* OR hypercholester?emia* OR hyperlipoprotein?emia* OR hypertriglycerid?emia*)
 # 9 TS=(“high blood pressure”)
 # 8 TS=(hypertensi* OR “peripheral arter* disease”)
 # 7 TS=(stroke OR stokes OR cerebrovasc* OR cerebral OR apoplexy OR (brain SAME accident*) OR (brain SAME infarct*))
 # 6 TS=(“atrial fibrillat*” OR tachycardi* OR endocardi*)
 # 5 TS=(pericard* OR isch?em* OR emboli* OR arrhythmi* OR thrombo*)
 # 4 TS=(cardio* OR cardia* OR heart* OR coronary* OR angina* OR ventric* OR myocard*)
 # 3 TS=(chitin* or lignin* or hemicellulose* or hexosane* or pentosane* or xanthan* or fructan* or inulin* or polyuronide* or pectin* or alginate* or “alginic acid*” or agar* or carrageen* or raffinose* or xylose* or polydextrose* or lactulose* or cellulose* or fibre* or fiber* or high-fibre* or high-fiber*)
 # 2 TS=(diet* or consum* or eat* or food* or nutri*)
 # 1 TS=(“dietary fiber” or “dietary fibre” or roughage* or prebiotic* or “carbohydrate polymer*” or non-starch or “non starch” or polysaccharide* or polysaccharide*)

CONTRIBUTIONS OF AUTHORS

All authors contributed to protocol development. The Trials Search Co-ordinator of Cochrane Heart ran the searches. LH, MM, and KR screened titles and abstracts and assessed studies for formal inclusion and exclusion. LH, MM, EL and JC abstracted data and assessed methodological rigour. Analyses were conducted by EL and checked by JC and KR. LH wrote the first draft of the results which was updated by EL and KR, and KR wrote the remaining sections.

DECLARATIONS OF INTEREST

LH - None

MM - My work in the pharmaceutical industry as a Regulatory Affairs Consultant is not related in any way to this review. I was not involved in development, design or any review of efficacy of drug products. My work is based only in chemistry manufacturing and controls to ensure quality of established drug products for license changes.

EL - None

JC - None

KR - None

SOURCES OF SUPPORT

Internal sources

- Warwick Medical School, University of Warwick, UK.

External sources

- National Institute for Health Research Cochrane Programme Grant, UK.
- Karen Rees is also supported by the National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care West Midlands at University Hospitals Birmingham NHS Foundation Trust, UK.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We planned to search Google Scholar, Open Grey, conduct a citation search and contact experts. There was insufficient time and resources to do this.

We excluded participants with type 2 diabetes. Whilst this is a major risk factor for CVD, interventions for the treatment and management of type 2 diabetes are covered by reviews registered with the Cochrane Metabolic and Endocrine Disorders Group.

We excluded trials focused on weight loss in order to avoid confounding.

We did not use clinical endpoints in subgroup analyses as these were not reported in included trials.